

Plants of Western Australian granite outcrops

S D Hopper¹, A P Brown² & N G Marchant³

¹Kings Park and Botanic Garden, West Perth WA 6005
email: steveh@kpbg.wa.gov.au

²Western Australian Threatened Species & Communities Unit,
Department of Conservation and Land Management, PO Box 51, Wanneroo WA 6065
email: andrewbr@calm.wa.gov.au

³Western Australian Herbarium, Department of Conservation and Land Management,
PO Box 104, Como WA 6152

Abstract

Outcropping granite rocks in Western Australia span a considerable climatic range, from the mediterranean south-west to inland desert and northern arid subtropics and tropics. At least 1320, and possibly 2000, plant taxa occur on Western Australian granite outcrops. Outcrop plant life is most diverse in the South West Botanical Province, with individual outcrops having up to 200 species, including many endemics not found in surrounding habitats. Species richness and local endemism declines with increasing aridity, to the point where Kimberley and Pilbara outcrops show little discontinuity in species from the surrounding landscape matrix. Outcrops are dominated by woody and herbaceous perennials, especially of the Myrtaceae, Orchidaceae, and Mimosaceae, and have an unusually rich diversity of annuals (Asteraceae, Stylidiaceae, Poaceae, Amaranthaceae etc.) compared with the flora as a whole. An unusual life form is found in resurrection plants capable of extreme desiccation and rehydration (e.g. *Borya*, *Cheilanthes*). Among woody perennials, bird pollination is frequent, and some outcrops harbour a high proportion of obligate seeder species due to the refuge from fire provided by bare rock barriers. The diversity of microhabitats and soil moisture regimes on outcrops has enabled the persistence of species beyond their main range in the face of climatic fluctuations. It has also facilitated the evolution of many endemics in the south-west. Major threatening processes facing outcrop plant communities include weed invasion, grazing by stock and feral animals, too-frequent fire, clearing, loss of shrub layer, salinity, and dieback. Conservation of these rich rare habitats needs the support of local communities.

Introduction

Western Australia is the largest Australian State, extending 2 391 km north-south and 1 621 km east-west, and occupying 2 525 500 km², or about a third of the nation. Climates range from temperate and mediterranean in the south-west to the monsoonal tropics in the extreme north, with arid conditions prevailing over much of the State from the Nullarbor Plain north through the inland deserts to the north-west Pilbara and Kimberley regions.

Most of the State is a plateau 300-600 m above sea level, with the highest point (Mt Meharry) only 1 251 m. Western Australia's numerous granite inselbergs and outcrops, consequently, are significant landscape features, even though most emerge less than 100 m above surrounding terrain.

Granite (or granitoid) rocks are mainly found on the Yilgarn and Pilbara Cratons, which together occupy about a third of the State's landmass, and along adjacent south coastal and Kimberley orogens (Myers 1997). The outcrops thus span a considerable climatic range, from the mediterranean south-west to inland desert and northern arid subtropics and tropics.

Western Australian outcrops vary in area from about

one tenth of a hectare to many hectares. While most are clearly isolated and disjunct, sometimes separated from the next outcrop by tens of kilometres, others are part of extensive inselberg systems covering several square kilometres (e.g. outcrop systems in Cape Le Grand, Porongurups and Cape Arid National Parks on the south coast, at Oudabunna Station near Paynes Find, and at Woodstock Station in the Pilbara). Peak Charles, 100 km north-west of Esperance, stands highest above surrounding country among the State's inselbergs, reaching 500 m from a low plain.

While the Kimberley, Pilbara and desert granites are in regions with depauperate floras not unlike those of much of tropical and arid Australia, the granite of the Yilgarn Craton underlies one of the world's regions richest in plant life, the South West Botanical Province (Marchant 1973; Hopper 1979, 1992; Beard 1980, 1981, 1990). As many as 8000 vascular plant species occur in this region, with about 75% endemism. Consequently, the botanical exploration of south-western inselbergs has been particularly rewarding, and much remains to be done.

Here, we provide an overview of current knowledge of plant life on Western Australian granite outcrops. Our aim was to compile a baseline, including a preliminary list of plants (vascular and cryptogam) collected from Western Australian outcrops, and a list of key references, to inform future research and provide a context for more specific studies. We conclude with some thoughts on the

conservation of this special component of Western Australian plant life.

Botanical Exploration

Western Australian granite outcrops have been important to aboriginal people ever since they occupied the land, but little has been documented of this long-standing relationship (Bindon 1997). Recorded European knowledge of granite outcrop plants in the State commenced with Archibald Menzies, surgeon and naturalist aboard the *HMS Discovery*, under the command of Captain George Vancouver. The *Discovery* was anchored in King George's Sound from September 28 to October 11, 1791.

Menzies made a "copious collection of ... vegetable productions, principally the genus *Banksia*, which are here very numerous" from various sites onshore in the vicinity of present-day Albany (Maiden 1909). Given the prominence of granite headlands, hills and islands at Albany, Menzies undoubtedly collected from them, but details are difficult to trace.

European botanists first named granite outcrop plants from Western Australia in 1800. The Frenchman Jacques J H de Labillardiere was naturalist on Admiral Bruny d'Entrecasteaux's expedition to Australia, which sought refuge from a storm near the modern town of Esperance over the period 9-17 December 1792. Labillardiere's specimens included a plant he named in 1800 as *Eucalyptus cornuta*, collected from the granite Observatory Island on 13 December 1792. *E. cornuta* occupies the apron and deeper soil pockets on granite outcrops throughout the Esperance area and adjacent islands, extending westwards to the Cape Naturaliste area (Fig 1).

Labillardiere named other granite outcrop plants from the expedition, including the bizarre genus *Borya* (Fig 4), a pincushion lily and resurrection plant, named in 1805 and typified by the species *B. nitida* which occurs on shallow soils on outcrops along the south coast from Albany to east of Esperance.

Robert Brown, on the expedition to circumnavigate Australia led by Matthew Flinders in the *Investigator*, anchored at King George's Sound on December 8, 1801, and collected extensively until January 5, 1802. Between January 10-18, the expedition also explored and collected near Esperance, landing at Lucky Bay (in the present-day Cape Le Grand National Park) for five days, and on islands in the Recherche Archipelago. Brown (1810) also named many granite outcrop plants, including another species of *Borya* (*B. spaherocephala*) that occurs with *B. nitida* on several south coast outcrops.

The arrival of James Drummond at the Swan River with Captain Stirling's colonising party on the *Parmelia* in 1829 heralded a major advance in knowledge of the south-west's flora, including that of granite outcrops (Erickson 1969). Drummond, a Scot and keen nurseryman with an excellent eye for new taxa, collected widely throughout the forests and present day wheatbelt, his specimens destined for subscribers in Great Britain and Europe. He was followed by many others up to modern times. Beard (1981) provides a useful historical review. The floral wealth of the south-west is such that significant numbers of new taxa

continue to be discovered and described each year (Green 1985; Hopper 1992), including granite outcrop endemics.

The ecology and biogeography of Western Australian granite outcrop plants were observed by the earliest collectors, but drew specific mention by the plant geographer Ludwig Diels (1906), who noted the "dwarflike" vegetation. Smith (1962) provided an account of the vegetation of the granite Porongurup Range, and many others have mentioned granite outcrop plants in broader vegetation surveys in the south-west (e.g. Beard 1981; Newbey & Hnatiuk 1985; Newbey *et al.* 1995; Wardell Johnson & Williams 1996; Brooker & Margules 1996).

Main (1967) gave a delightful account of the natural history of Yorkrakine Rock through an annual cycle. Marchant (in Erickson *et al.* 1973) summarised botanical features of south-western outcrops in an essay for a general readership.

Schweinfurth (1978) described the vegetation of exposed granite headlands along the south coast, while Abbott & Watson (1978) and Abbott (1980) explored these headlands and adjacent islands in greater detail, showing the flora of most islands had a depauperate subset of that found onshore, and displayed little endemicity nor genetic differentiation from mainland populations.

Hopper (1981) highlighted the importance of winter-flowering woody perennials as a nectar resource used by a diversity of honeyeaters in the central wheatbelt (Fig 3). Pate & Dixon (1982) gave a comprehensive account of the biology of tuberous and cormous plants, including many found on granite.

Ornduff (1987) documented the flora of herbfields including, and centrewards from, the *Borya* zone on nine outcrops near Perth and Hyden. Burgman (1987) sampled five outcrops inland from Esperance and explored aspects of rarity in granite communities. Hopper *et al.* (1990) illustrated 29 endangered endemics of granite outcrops and highlighted conservation issues. Pignatti & Pignatti (1994) reported on herbfields from shallow seasonally wet soils on a selection of south-west outcrops. Ohlemüller (1997) documented plants in shallow soil depressions on 26 outcrops throughout the south-west and goldfields.

In parallel with these ecological investigations, genetic studies of Western Australian granite outcrop plants were pioneered by James (1965), who's penetrating work on the herb *Isotoma petraea* (Lobeliaceae) has become a classic in the literature on plant evolution (Bussell & James 1997). Studies on the population genetics of endemic granite outcrop eucalypts have also provided useful insights into the evolution of highly fragmented small populations (Moran and Hopper 1983; Sampson *et al.* 1988). The breeding systems of granite outcrop endemics have been documented for *Villarsia* (Menyanthaceae; Ornduff 1986, 1996) and *Anthocercis gracilis* (Solanaceae; Stace 1995).

Sampling Methods

Rather than laboriously search the increasing number of botanical publications that include plant taxa recorded

on Western Australian granite outcrops, we compiled the plant list from specimens housed in the Western Australian Herbarium. The database WAHERB stores full label details of 413 000 specimens of vascular plants and cryptogams collected throughout the State, and afforded an opportunity to list those taxa for which habitat details on labels included the words "granite outcrop". Initial trials indicated that searching on "granite" or "granitic" alone picked up too many taxa that occupied regions underlain by granite rock, but favoured deep soils well removed from rock outcrops.

In our list, taxa are mainly species, but infraspecific subspecies, varieties and forms, together with hybrids, are all counted. This ensures that all named plant biodiversity is included, and allows for discrepancies in assigning rank among taxonomists.

The above approach to listing plant taxa was based on an understanding that most authors involved in botanical survey of Western Australian granite outcrops have collected voucher specimens and deposited them in the Western Australian Herbarium (e.g. Ornduff 1987; Burgman 1987; Hopper 1981; Hopper *et al.* 1990; Hopper & Brown, *unpublished orchid research*; Newbey & Hnatiuk 1985; Newbey *et al.* 1995). In addition, as an independent check, the senior author has compiled field herbaria in notebooks for *ca* 150 outcrops throughout Western Australia over the past two decades. Most outcrops sampled occur in the South West Botanical Province (e.g. Fig 1) and adjacent pastoral region, while a few have been examined in the Pilbara (e.g. Fig 2) and Kimberley.

Precisely defining the boundaries of a granite outcrop is difficult, especially for mobile components of the biota (Main 1997). However, the sedentary nature of adult plants, especially perennials, affords a reliable and measurable biological indicator of the limits of the outcrop system.

In fieldwork, we have used the distribution of plants that are locally endemic to outcrops to define the extent of outcrop communities sampled. Thus, all species centrewards from the outermost individuals of granite outcrop local endemics were included as present on an outcrop. This approach ensures that all endemics are sampled, but has the effect of picking up some species from surrounding deeper soils that do not extend onto soil pockets on the outcrop itself. Other authors have restricted their sampling to such soil pockets (e.g. Houle 1990; Porembski *et al.* 1995; Ohlemuller 1997), or only to herbfields (e.g. Ornduff 1987). Experience showed that perennial vegetation fringing the base of Western Australian outcrops had to be sampled to ensure a comprehensive inventory (see below).

We followed Newbey and Hnatiuk (1985) and Newbey *et al.* (1995) in collecting on random walks stratified by microhabitat rather than within quadrats *i.e.* the *random stratified walk* technique. A comparison of the senior author's data with the quadrat-based data of Burgman (1987) for Mt Ney inland from Esperance showed that four 5m x 5m quadrats spaced at 25 m intervals on NE scree slope and sheet rock yielded 61 taxa, whereas 192 taxa were documented through random stratified walks covering cardinal compass points.

Microhabitats recognised and deliberately searched

for on random stratified walks included bare rock, cryptogamic crusts, gnammas (rock pools or weather pits, seasonal and permanent), soil-filled crevices, caves/tafoni/shade of boulders or exfoliated slabs, herbfields on shallow well-drained soil, herbfields on shallow seasonally waterlogged soil (ephemeral flush vegetation), shrublands and woodlands on deep well-drained soil, shrublands, woodlands and forests on deep seasonally waterlogged soil, permanent springs, major watercourses, and salt lakes.

Floristics

A total of 1320 Western Australian granite outcrop plant taxa were represented and labelled as such in the collections of the Western Australian Herbarium in May 1996 (Appendix 1). Of these, 1097 taxa (83.0%) were described, 94 (7.1%) were undescribed but well-delimited with manuscript names, and 129 (9.9%) were of uncertain taxonomic delimitation. Only 4 (0.3%) were hybrids, well below the proportion (4%) estimated for the whole WA flora (Hopper 1995). The list includes 11 macrofungi, 34 lichens, 46 mosses, 9 liverworts, 9 ferns, 7 fern allies, 2 cycads, 2 conifers, and 1200 angiosperms (284 monocotyledons, 916 dicotyledons).

While these statistics provide a useful baseline, it is pertinent to note that sampling of most cryptogam groups is poor. For example, we recorded only 34 lichens with vouchered granite outcrop specimens for the whole of WA, whereas Pigott & Sage (1997) list 36 lichen taxa for Yilliminning Rock alone. Macrofungi undoubtedly are orders of magnitude more diverse than presently known. The moss flora is comparatively better documented at the species level (Stoneburner *et al.* 1993; Stoneburner & Wyatt 1996), but much more collecting is needed to fill in knowledge of distribution and ecology.

Some confidence may be placed in the list of vascular plant taxa, which accounts for 1220 (92.4%) of the total of 1320. Based on our knowledge of rock outcrop floras across the State, the majority of south-western vascular plant taxa on granite outcrops are listed. However, some conspicuous omissions of taxa that are common (e.g. *Hakea petiolaris*, *Corymbia calophylla*), uncommon (e.g. *Grevillea magnifica*, *Phylloglossum drummondii* and *Isoetes drummondii*) and rare (e.g. *Drummondita hassellii* var *longifolia*, *Hibbertia bracteosa*, *Villarsia calthifolia*, *Pleurophascum occidentale*) indicate that revision upwards is required. Moreover, a comparison of the list of 187 taxa for Yilliminning Rock near Narrogin (Pigott & Sage 1997) with Appendix 1 shows that 78 Yilliminning taxa (42%) are not present in our list for the State. This discrepancy would be even more so for northern outcrops. For example, absent from the present list are the dominant spinifex hummock grasses on granite outcrops (*Triodia* spp), as well as common lemon grasses (*Cymbopogon* spp), and scattered but widespread subtropical woody shrubs and small trees such as *Terminalia canescens*. It would be no surprise to us, consequently, if the number of vascular plant taxa on Western Australian granite outcrops approached 2000 in the future.

In the present list, major vascular plant families include the Myrtaceae (162 taxa), Orchidaceae (159), Mimosaceae

(82), Asteraceae (72), Papilionaceae (66), Proteaceae (54), and Stylidiaceae (33). Genera richest in taxa on granite outcrops include *Acacia* (82 taxa), *Caladenia* (70), *Eucalyptus* (44), *Stylium* (28), *Melaleuca* (26), *Grevillea* (25), *Drosera* (21), *Verticordia* (20), *Eremophila* (18) and *Thelymitra* (16).

Thus, woody perennial shrubs and trees of myrtles, wattles, peas and Proteaceae are taxon-rich on WA granite outcrops, reflecting trends in the flora as a whole (Green 1985; Hopper *et al.* 1996). However, the granite woody perennials are depauperate in Epacridaceae, which is rich in other habitats. The granite outcrop flora is clearly divergent from the flora at large in the unusually high taxon richness of groups such as the tuberous perennial terrestrials (orchids, sundews, lilies), and herbaceous, often annual, daisies and triggerplants.

Systematic sampling of shallow-soil herbfields on south-western outcrops by Ornduff (1987) and Ohlemüller (1997) yields a useful comparison with our data on herbaceous families. Ornduff recorded 160 taxa belonging to 37 families (he didn't sample Orchidaceae for conservation reasons), of which the largest were Asteraceae, Poaceae, Liliaceae *sens. lat.*, and Stylidiaceae. Ohlemüller recorded 134 species from 42 families, with the largest being Asteraceae, Orchidaceae, Poaceae, Apiaceae, Stylidiaceae and Centrolepidaceae.

Major herbaceous families in our list are Orchidaceae (Fig 5), Asteraceae, Stylidiaceae, Anthericaceae (or Liliaceae *sens. lat.*), Poaceae, Droseraceae, Cyperaceae, Goodeniaceae, Amaranthaceae and Centrolepidaceae. The Amaranthaceae is a family richest in the north of Western Australia, so it is not surprising that Ornduff (1987) and Ohlemüller (1997) found it to be depauperate in their south-west study areas. Orchidaceae undoubtedly emerges as the most taxon-rich herbaceous family in our list because two of us (SH and AB) have made a detailed and prolonged study of granite outcrop orchids, and found that repeat surveys over different seasons and years are needed to develop comprehensive orchid inventories for individual rocks. Other differences between our ranking and those of Ornduff (1987) and Ohlemüller (1997) are minor. This general concordance suggests that major trends and statistics for the vascular plants in Appendix 1 are accurate, at least for south-west outcrops.

Plants of Special Interest

Granite outcrop habitats have elicited remarkable parallel evolution in many plants and animals (Porembski *et al.* 1997; Wyatt 1997; Mares 1997). The combination of high solar radiation, rapid runoff of rainfall, and shallow soils on a rocky substrate provide microhabitats of accentuated seasonal and diurnal stresses. Conditions may vary over a few metres from cool permanently moist shaded caves with water seepage to drought-afflicted shallow soils and rock surfaces fully exposed to all the elements.

Few organisms can tolerate the harshest of these rock-surface habitats, which tend to be occupied by cryptogamic crusts of cosmopolitan cyanobacteria, lichens and mosses such as *Grimmia laevigata*. Western Australian outcrops conform to this trend. However, on

south-west outcrops experiencing high rainfall, moss cushions become prominent, with the luxuriant brown carpets of *Bretelia affinis* and the verdant green *Campylopus bicolor* and *C. australis* noteworthy.


Coping with seasonal or unpredictable drought is, undoubtedly, the most significant survival strategy faced by granite outcrop plants. Among Western Australian herbaceous perennials on outcrops, pincushion lilies (*Borya* spp, Boryaceae) are abundant and remarkable in their capacity as resurrection plants (Gaff 1981) to withstand desiccation to less than 5% of normal leaf moisture content, turning orange in the process, and rehydrating to normal green leaves within a day of rainfall (Fig 4). There are at least six species of *Borya* in the South West Botanical Province (*B. sphaerocephala* is a variable taxon awaiting detailed study and possible subdivision). These plants do not extend to the Pilbara outcrops. Other resurrection plants commonly seen on WA granites include the rock ferns *Cheilanthes* spp and *Pleurozorus rutifolius*.

Persisting underground as a tuber during drought is a common strategy among Western Australian granite outcrop herbs in the south-west and adjacent pastoral regions, especially lilioids (e.g. *Wurmbea* spp, Colchicaceae), orchids (*Caladenia*, *Thelymitra*, *Pterostylis*, *Diuris*, *Prasophyllum*) and sundews (*Drosera*), as well as the unusual fern ally *Phylloglossum* (Pate & Dixon 1982). On northern outcrops, tuberous Cyperaceae tend to occupy this niche.


Succulence is not prominent in the Western Australian granite outcrop perennial floras, but has evolved in a few taxonomically-unrelated species e.g. *Spiculaea ciliata* (Orchidaceae; Fig 5), *Carpobrotus* spp (Aizoaceae). Some perennial shrubs of south coast granites have unusually thick leaves (e.g. *Hakea clavata*, Proteaceae; *Anthocercis viscosa*, Solanaceae), and at least one has tuberous roots (*Calothamnus tuberosus*, Myrtaceae). Sclerophyllous leaves characterise the vast majority of woody perennials found on granite outcrops, as well as in the Western Australian flora at large.

The graminoid habit has evolved in several unrelated groups in the south-west, with common perennials forming tussocks including diverse Cyperaceae (especially *Lepidosperma* spp), blind grass (*Stypandra* spp, Phormiaceae), the sedge-like grass *Spartochloa scirpoidea* (Poaceae) and lemon grass *Cymbopogon* spp (Poaceae). Desert, Pilbara and Kimberley outcrops are dominated by uniquely Australian hummock grasses with inrolled pungent leaves (*Triodia* spp, Poaceae).


Drought avoidance through an annual life history is a major feature of Western Australian granite herbs in families such as the Asteraceae, Stylidiaceae, Poaceae, Goodeniaceae, Amaranthaceae and Centrolepidaceae. Most of these annuals have relatively small inconspicuous flowers suggestive of inbreeding (Ornduff 1987), but some are brightly coloured and abundant. Most outcrops in southern Western Australia are bedecked with colourful swards of annual triggerplants (*Stylium*, Stylidiaceae) and everlasting (e.g. *Rhodanthe*, *Waitzia*, Asteraceae) that enliven the herbfields. Succulence is a feature of some annuals, especially those that occupy the dry spectrum of shallow soils (e.g. *Calandrinia* spp, Portulacaceae; *Crassula* spp, Crassulaceae).




1. Coastal granite outcrops east of Albany on Two Peoples Bay Nature Reserve, with *Eucalyptus cornuta* (pale canopy), described in 1800 by French scientific explorer Labillardiere. Photograph by SD Hopper.




2. Granite sheet, boulders and distant inselbergs in the arid Pilbara, Spear Hill, south-west of Marble Bar. Photograph by SD Hopper.




3. Bird-pollinated plants are a feature of south-west Australian granite outcrops - New Holland Honeyeater on *Eucalyptus caesia*, Boyagin Rock, north-west of Pingelly. Photograph by SD Hopper.



4. The resurrection plant, pincushion lily *Borya constricta*, with leaves 1 cm long desiccated (orange) but rehydrating (green) following rain at the end of summer drought, Chiddarcooping Rock, north-east of Merredin. Photograph by SD Hopper.



5. Terrestrial orchids are the second most species-rich plant family on Western Australian granite outcrops. Here, flowers of the elbow orchid (*Spiculaea ciliata*) 1 cm long splay from the succulent stem that feeds them in early summer, and a male thynnid wasp attempts to copulate with the flower's sexual decoy, an insectiform hinged labellum, Boulder Rock, Brookton Highway. Photograph by BA & AG Wells.



6. Arguably the rarest Western Australian granite outcrop plant, the annual aquatic millfoil *Myriophyllum lapidicola* in full flower (with round floating leaves), and known from just two gnammas (rock pools), alongside the invasive aquatic South African weed, *Crassula natans* var *minus*. Photograph by SD Hopper.

Annuals are also found commonly in seasonally waterlogged shallow soils or ephemeral flush communities (Pignatti & Pignatti 1994). These annuals include species of Centrolepidaceae (*Centrolepis* and *Aphelia*), Apiaceae (*Hydrocotyle*), Juncaginaceae (*Triglochin*), Cyperaceae (*Schoenus*, *Isolepis*, *Cyperus*), Asteraceae (*Quinetia*, *Millotia*, *Rutidosis*, *Siloxerus*, *Toxanthes*, *Hyalosperma*, *Podotheca* etc.) and Styliadiaceae (*Styliodium*, *Levenhookia*).

Deeper soils enable survivorship of woody perennials, of which the largest on Western Australian outcrops include eucalypts (*Eucalyptus* and *Corymbia*, Myrtaceae), wattles (*Acacia*, Mimosaceae), she-oaks (*Allocasuarina*, Casuarinaceae), and rock figs (*Ficus*, Moraceae). A noteworthy feature of the woody perennials is the high proportion of bird-pollinated species (Hopper 1981; Fig 3) in families such as Myrtaceae (e.g. *Eucalyptus*, *Calothamnus*, *Melaeuca*, *Verticordia*), Proteaceae (*Grevillea*, *Hakea*, *Banksia*), Myoporaceae (*Eremophila*), Papilionaceae (*Kennedyia*, *Crotalaria*), Viscaceae (*Amyema*, *Lysiana*) and Sterculiaceae (*Brachychiton*). Also, south-western outcrops have an unusually high number of woody perennials that are obligate seeders - plants that are killed by fire and recruit only from seed. For example, 77% of the perennials in a granite community at Chiddarcooping Nature Reserve north-east of Merredin were obligate seeders (Hopper *et al.*, *unpubl.*).

Gnammas on Western Australian outcrops have a unique flora comprising annual species of quillworts (*Isoetes*, Isoetaceae), mudmats (*Glossostigma*, Scrophulariaceae), millfoils (*Myriophyllum*, Haloragaceae; Fig 6) and others, including the introduced South African annual *Crassula natans* (Crassulaceae).

Weeds

Ornduff (1987) and Ohlemüller (1997) found that introduced weed species, mainly annuals, comprised 23.7% and 17.0% of the granite herbfield floras sampled respectively. These figures are high compared with the ca 9% that weeds represent of the WA flora as a whole (Green 1985; Keighery 1995). They are even more significant when considered with granite outcrop herbfield floras elsewhere on earth, which have far fewer invasive weeds (Porembski *et al.* 1997; Wyatt 1997).

Weed growth is most pronounced in full sun on outcrops, especially where soil has been disturbed and enriched by rabbit dung or agricultural activity. In these situations, annual grasses such as *Briza maxima*, *Avena fatua* and *Ehrharta longifolia* often dominate and replace native annuals throughout much of the south-west. Weeds are rare only where a dense shrub layer or low forest of native woody perennials persist. It is clear that a persistent seed rain of weeds occurs over vast regions of the south-west, as weeds appear in open areas on outcrops where little or no disturbance is evident and native plants dominate.

Hopper (1997) has attributed the high invasibility of disturbed Western Australian plant communities to the absence of major glacial soil stripping as an evolutionary force acting on the flora. Native species are unable to compete against weeds from habitats where soil disturbance is a regular perturbation. Further

experimental study of weeds in granite herbfields could test this hypothesis, and assist attempts at restoration of invaded outcrops.

Biogeography and Endemism

Western Australian granite outcrops display plant biogeographic patterns that mirror that of the whole flora (Hopper 1979, 1992; Hopper *et al.* 1996); species-richness and endemism are pronounced in the transitional rainfall zone (wheatbelt) of the south-west, and attenuate as rainfall decreases through the pastoral country to the deserts, Pilbara and Kimberley. For example, the total number of vascular plant taxa recorded by the senior author and colleagues on a sample of rocks ranged from 142 (Point Matthew, near Augusta) and 201 (Mt Frankland) in the highest rainfall forests, 192 (Mt Ney) and 187 (Yilliminning Rock near Narrogin; Pigott and Sage 1997) in the transitional rainfall zone, to 85 (Daggar Hills, near Yalgoo) in the pastoral zone, and 90 (Mooleyella Rocks, east of Marble Bar) and 80 (Spear Hill, west of Marble Bar; Fig 2) in the arid Pilbara.

There are no vascular species shared between northern (Kimberley, Pilbara) outcrops and those in the south-west. Moreover, the northern outcrop floras are virtually identical with that from the matrix of surrounding terrain, save for outcrop specialists like rock figs and rock ferns. Walters & Wyatt (1982) similarly recorded low endemism and little discontinuity between vascular plants on granite outcrops and adjacent landforms of the arid Central Mineral Region of Texas.

In contrast, south-western and adjacent pastoral zone rocks have higher levels of local endemism, especially in the high rainfall forest region where the outcrops present the most striking difference in habitat to the surrounding vegetation matrix (e.g. Wardell Johnson & Williams 1996; Brooker & Margules 1996).

Biogeographical relationships of outcrop floras across the south-west are under ongoing study by the senior author. As a precursor, Hopper & Brown (*unpublished*) documented the distribution of 126 orchid taxa on 41 outcrops ranging from the highest rainfall forests through the transitional zone wheatbelt to the arid zone. Each rock outcrop was treated as a site in a classification of the orchid data.

The study highlighted a number of significant trends. A primary division occurred between the 15 rocks found in the forested High Rainfall Zone (>800 mm p.a.) and the rest ranging from the Transitional Rainfall Zone of the wheatbelt into the Arid Zone. Subsequent divisions established as much difference among forest rocks as among the wheatbelt/arid rocks, even though the forest rocks were confined to a much smaller area. Moreover, remarkably, closely adjacent rocks were widely separated in the classification, indicating significant differences in their orchids (e.g. 10.3 km Rock and 10.9 km Rock of Ornduff's (1987), separated by just 600 m of jarrah forest on Albany Highway). Conversely, rocks separated geographically often had similar orchid floras (e.g. Boyagin Rock and Pingaring Rock on the western and eastern sides of the south-central wheatbelt respectively).

These patterns suggest significant barriers to orchid

Table 1
Lists of Western Australian orchid taxa that occur on granite outcrops.

All taxa		
<i>Caladenia attingens</i> subsp. <i>attingen</i> ms	<i>Caladenia voigtii</i> ms	<i>Pterostylis aspera</i>
<i>Caladenia attingens</i> subsp. <i>gracillima</i> ms	<i>Corybas recurvus</i>	<i>Pterostylis barbata</i>
<i>Caladenia brevisura</i> ms	<i>Cyanicula amplexans</i> ms	<i>Pterostylis elegantissima</i>
<i>Caladenia brownii</i> ms	<i>Cyanicula ashbyae</i> ms	<i>Pterostylis hamiltonii</i>
<i>Caladenia caesarea</i> subsp. <i>maritima</i> ms	<i>Cyanicula caerulea</i> subsp. <i>apertala</i> ms	<i>Pterostylis mutica</i>
<i>Caladenia caesarea</i> subsp. <i>transiens</i> ms	<i>Cyanicula deformis</i> ms	<i>Pterostylis recurva</i>
<i>Caladenia citrina</i> ms	<i>Cyanicula fragrans</i> ms	<i>Pterostylis roensis</i>
<i>Caladenia denticulata</i>	<i>Cyanicula gemmata</i> ms	<i>Pterostylis sanguinea</i>
<i>Caladenia dimidia</i> ms	<i>Cyanicula sericea</i> ms	<i>Pterostylis sargentii</i>
<i>Caladenia discoidea</i>	<i>Cyrtostylis huegelii</i>	<i>Pterostylis scabra</i>
<i>Caladenia doutchiae</i>	<i>Cyrtostylis robusta</i>	<i>Pterostylis vittata</i>
<i>Caladenia extans</i> ms	<i>Diuris</i> aff. <i>longifolia</i>	<i>Pyrorchis nigricans</i>
<i>Caladenia falcata</i>	<i>Diuris brumalis</i>	<i>Spiculaea ciliata</i>
<i>Caladenia filifera</i>	<i>Diuris conspicillata</i>	<i>Thelymitra</i> aff. <i>holmsii</i>
<i>Caladenia flaccida</i> subsp. <i>flaccida</i> ms	<i>Diuris laevis</i>	<i>Thelymitra</i> aff. <i>longifolia</i>
<i>Caladenia flaccida</i> subsp. <i>pulchra</i> ms	<i>Diuris laxiflora</i>	<i>Thelymitra</i> aff. <i>nuda</i>
<i>Caladenia flava</i> subsp. <i>flava</i> ms	<i>Diuris longifolia</i>	<i>Thelymitra</i> aff. <i>pauciflora</i>
<i>Caladenia flava</i> subsp. <i>maculata</i> ms	<i>Diuris maculata</i>	<i>Thelymitra</i> antennifera
<i>Caladenia flava</i> subsp. <i>sylvestris</i> ms	<i>Diuris picta</i>	<i>Thelymitra</i> benthamiana
<i>Caladenia footeana</i> ms	<i>Diuris pulchella</i>	<i>Thelymitra</i> crinita
<i>Caladenia granitora</i> ms	<i>Diuris recurva</i>	<i>Thelymitra</i> cucullata
<i>Caladenia heberleana</i> ms	<i>Drakonorchis barbarossa</i> ms	<i>Thelymitra</i> aff. <i>dedmaniarum</i>
<i>Caladenia hirta</i> subsp. <i>rosea</i> ms	<i>Drakonorchis drakeoides</i> ms	<i>Thelymitra</i> flexuosa
<i>Caladenia hoffmannii</i> subsp. <i>graniticola</i> ms	<i>Drakonorchis mesocera</i> ms	<i>Thelymitra</i> macrophylla
<i>Caladenia incensa</i> ms	<i>Elythranthera brunonis</i>	<i>Thelymitra spiralis</i>
<i>Caladenia incrassata</i> ms	<i>Elythranthera emarginata</i>	Taxa that predominantly occur on granite outcrops
<i>Caladenia infundibularis</i>	<i>Eriochilus dilatatus</i> subsp. <i>dilatatus</i> ms	<i>Caladenia caesarea</i> subsp. <i>maritima</i> ms
<i>Caladenia integra</i>	<i>Eriochilus dilatatus</i> subsp. <i>multiflorus</i> ms	<i>Caladenia extans</i> ms
<i>Caladenia latifolia</i>	<i>Eriochilus dilatatus</i> subsp. <i>undulatus</i> ms	<i>Caladenia granitora</i> ms
<i>Caladenia lobata</i>	<i>Eriochilus helonomos</i> ms	<i>Caladenia hoffmannii</i> subsp. <i>graniticola</i> ms
<i>Caladenia longicauda</i> subsp. <i>clivicola</i> ms	<i>Eriochilus pulchellus</i> ms	<i>Caladenia integra</i>
<i>Caladenia longicauda</i> subsp. <i>eminens</i> ms	<i>Eriochilus scaber</i>	<i>Caladenia longicauda</i> subsp. <i>clivicola</i> ms
<i>Caladenia longicauda</i> subsp. <i>longicauda</i> ms	<i>Genoplesium nigricans</i>	<i>Caladenia longicauda</i> subsp. <i>rigidula</i> ms
<i>Caladenia longicauda</i> subsp. <i>rigidula</i> ms	<i>Leptoceras menziesii</i>	<i>Caladenia multiclavia</i>
<i>Caladenia longilabata</i>	<i>Lyperanthus serratus</i>	<i>Caladenia nivalis</i>
<i>Caladenia macrostylis</i>	<i>Microtis</i> aff. <i>parviflora</i>	<i>Caladenia remota</i> subsp. <i>remota</i> ms
<i>Caladenia marginata</i>	<i>Microtis atrata</i>	<i>Cyanicula ashbyae</i> ms
<i>Caladenia microchila</i> ms	<i>Microtis brownii</i>	<i>Cyanicula fragrans</i> ms
<i>Caladenia multiclavia</i>	<i>Microtis eremaea</i>	<i>Diuris conspicillata</i>
<i>Caladenia nivalis</i> ms	<i>Microtis graniticola</i>	<i>Diuris picta</i>
<i>Caladenia pachychila</i> ms	<i>Microtis media</i> subsp. <i>eremicola</i>	<i>Diuris pulchella</i>
<i>Caladenia pholcoidea</i> ms	<i>Microtis media</i> subsp. <i>media</i>	<i>Eriochilus pulchellus</i> ms
<i>Caladenia polychroma</i> ms	<i>Monadenia bracteata</i>	<i>Microtis eremaea</i>
<i>Caladenia radialis</i>	<i>Paracaleana nigrita</i>	<i>Microtis graniticola</i>
<i>Caladenia reptans</i> subsp. <i>impensa</i> ms	<i>Paracaleana triens</i> ms	<i>Microtis media</i> subsp. <i>eremicola</i>
<i>Caladenia reptans</i> subsp. <i>reptans</i> ms	<i>Prasophyllum</i> aff. <i>parvifolium</i>	<i>Spiculaea ciliata</i>
<i>Caladenia rhomboidiformis</i>	<i>Prasophyllum brownii</i>	<i>Thelymitra</i> aff. <i>nuda</i>
<i>Caladenia roei</i>	<i>Prasophyllum cucullatum</i>	<i>Thelymitra</i> aff. <i>dedmaniarum</i>
<i>Caladenia saccharata</i>	<i>Prasophyllum elatum</i>	Declared Rare orchid taxa of granite outcrops
<i>Caladenia serotina</i> ms	<i>Prasophyllum fimbria</i>	<i>Caladenia caesarea</i> subsp. <i>maritima</i> ms
<i>Caladenia sigmaeidea</i>	<i>Prasophyllum gibbosum</i>	<i>Caladenia extans</i> ms
<i>Caladenia splendens</i> ms	<i>Prasophyllum gracile</i>	<i>Caladenia hoffmannii</i> subsp. <i>graniticola</i> ms
<i>Caladenia hiemalis</i> ms	<i>Prasophyllum parvifolium</i>	<i>Caladenia voigtii</i> ms
<i>Caladenia horistes</i> ms	<i>Prasophyllum ringens</i>	<i>Thelymitra</i> aff. <i>dedmaniarum</i>
<i>Caladenia pendens</i> subsp. <i>pendens</i> ms	<i>Pterostylis</i> aff. <i>nana</i>	
<i>Caladenia remota</i> subsp. <i>remota</i> ms	<i>Pterostylis</i> aff. <i>rufa</i>	
<i>Caladenia pendens</i> subsp. <i>talbotii</i> ms	<i>Pterostylis allantoidea</i>	

dispersal, particularly between forest rocks, and high levels of local extinction and stochastic events underlying the presence of orchids on individual rocks in the southwest and adjacent arid zone. There have been dynamic climatic fluctuations across the south-west for several million years as Australia drifted northwards and arid conditions overtook much of central Australia (Hopper

1979; Hopper *et al.* 1996). The diversity of microhabitats on granite outcrops provided refuge for plants adapted to both dry or wet conditions as the surrounding matrix waxed and waned climatically (Marchant 1973; Main 1997). Survivorship in small populations on granite refuges undoubtedly was a matter of chance in the face of such repeated climatic turmoil.

Interestingly, the above conditions of small disjunct outcrop populations undergoing recurrent stresses is predicted as ideal for genetic divergence and speciation (Grant 1981). Is this prediction borne out by studies of Western Australian outcrop plants? Table 1 provides a recently updated list of 141 orchid taxa recorded from Western Australian granite outcrops. Of these, 22 (16%) are more or less endemic. The endemics have geographical ranges from widespread on outcrops throughout the south-west (e.g. *Spiculaea ciliata*; Fig 5) to highly restricted to a few adjacent outcrops less than 10 km apart (e.g. *Caladenia caesarea* subsp. *maritima*, *Thelymitra* aff. *dedmaniarum*).

In terms of evolutionary origins, these endemics display at least three patterns (Hopper and Brown, unpublished):

- relictual, with no obvious close relatives and therefore likely to have been on granites for a long period of time (e.g. *Spiculaea ciliata*, a monotypic genus);
- derived by speciation from allopatric congeners of habitats other than granite (e.g. *C. granitaria*, from coastal granites east of Albany, sister species to *C. infundibularis* of western high rainfall forests and coastal heaths; *C. hoffmannii* subsp. *graniticola*, of east central wheatbelt outcrops, sister to *C. hoffmannii* subsp. *hoffmannii* of lateritic loams well to the northwest in the Northampton region); and
- derived by speciation from allopatric congeners of other granite outcrops (e.g. *C. extans*, from outcrops east of Esperance, sister to *C. integra* of western wheatbelt outcrops).

Thus, the orchid data do indeed support the hypothesis that conditions on south-west granite outcrops have facilitated genetic divergence and speciation. Genetic studies of a few other granite taxa lend further support, e.g. the herb *Isotoma petraea* (Bussell & James 1997), and eucalypts endemic to granite (Hopper & Burgman 1983; Sampson et al. 1988). Clearly, more work along these lines is needed.

There is a large number of granite endemics in south-western Australia, especially among the perennials that dominate the woody vegetation and herbfields. We have already shown that 16% of orchids on outcrops are endemic. For eucalypts, around 24% are endemic (Hopper, unpublished). The level of endemism for the whole granite outcrop flora is difficult to determine without more penetrating research, but there is no doubt that south-western Australia has higher levels than any other system documented (e.g. Walters & Wyatt 1982; Porembski et al. 1995, 1997).

The refugial opportunities offered by south-west granite outcrops are also evident in species that have highly disjunct outliers well removed from the main geographical distribution. These include populations on wet outcrop sites in much lower rainfall areas than the main species' stand (e.g. the Jilakin Rock stand of jarrah *Eucalyptus marginata*, the Twine Rock stand of *E. wandoo*, and the Kuendar stand of *E. rufa*). Conversely, arid-adapted species penetrate high rainfall areas on dry north-facing slopes of granites (e.g. populations of *Eucalyptus drummondii* west of Margaret River).

Conservation

Granite outcrops occupy a very small proportion of most Western Australian landscapes in which they occur. Especially in the south-west, the outcrop plant communities are, therefore, by definition, rare, and likely to contain rare species.

Hopper et al. (1990) found that endangered granite outcrop plants numbered 29 (12.2%) of the 238 plants declared in 1989 as Rare Flora under the Wildlife Conservation Act. These endangered plants ranged from large mallees (e.g. *Eucalyptus crucis* subsp. *crucis*) and small trees (*Acacia denticulosa*, *Banksia verticillata*), through compact shrubs (e.g. *Drummondia hasselii* var. *longifolia*, *Verticordia staminea*) and climbers (*Kennedia beckxiana*, *K. macrophylla*) to diminutive herbs (*Tribonanthes purpurea*) and annual aquatics (*Myriophyllum petraeum*). While some endangered taxa have several populations spread across a number of disjunct outcrops, some are confined to very few localities (e.g. *Myriophyllum lapidicola*, known from just two gnammas). Accidental destruction of such populations could be catastrophic. Conservation in the wild in such cases needs to be backed up by off-site activities such as germplasm storage and artificial propagation (underway at Kings Park and Botanic Garden for *M. lapidicola* and other critically endangered granite endemics; Dixon 1994).

Apart from endangered species, the diverse communities on south-western granite outcrops are noteworthy in the rapidity with which they change within and between rocks. They are complex, ever-changing, and rare in their own right.

While many granite outcrops have been spared direct clearing due to their unsuitability for agriculture, and some are included within conservation reserves, most face threatening processes that need management if the native biota is to persist. Such processes include replacement or damage by invasive weeds, feral animals, grazing, inappropriate fire regimes, clearing, loss of shrub layer, salinity and dieback disease. We have briefly addressed the issue of weeds above, and highlighted the importance of maintaining undisturbed soil and dense shrub layers to control weeds effectively. Such restoration activities require an ongoing presence and commitment to a given outcrop. Local communities are vital in this context.

Western Australians are fortunate in being custodians of a unique and diverse suite of granite outcrop plants. We hope that this brief review will stimulate others to study and conserve what is a remarkable heritage.

Acknowledgements: To all those colleagues who helped with field work, discussion and ideas, our sincere thanks. L Sage assisted with the analysis of granite orchid biogeography. R Wyatt and R Ornduff hosted SH on sabbatical in the USA in 1990, and greatly facilitated development of an international perspective on granite outcrop plants. S Porembski has recently broadened this international perspective. We are grateful for their interest and ideas.

References

- Abbott I 1980 The transition from mainland to island, illustrated by the flora and landbird fauna of headlands, peninsulas and islands near Albany, Western Australia. Journal of the Royal Society of Western Australia 63:79-92.

- Abbott I & Watson JR 1978 The soils, flora, vegetation and vertebrate fauna of Chatham Island, Western Australia. Journal of the Royal Society of Western Australia 60:65-70.
- Beard JS 1980 A new phytogeographic map of Western Australia. Western Australian Herbarium Research Notes 3:37-58.
- Beard JS 1981 Vegetation of Western Australia. 1:1 000 000 Series. Swan. University of Western Australia, Perth.
- Beard JS 1990 Plant Life of Western Australia. Kangaroo Press, Sydney.
- Bindon P 1997 Aboriginal people and granite domes. Journal of the Royal Society of Western Australia 80:173-179.
- Brooker MG & Margules CR 1996 The relative conservation value of remnant patches of native vegetation in the wheatbelt of Western Australia: I. Plant diversity. Pacific Conservation Biology 2:268-278.
- Brown R 1810 Prodromus Florae Novae Hollandiae et Insulae Van-Diemen exhibens characteres plantarum quas annis 1802-1805. Taylor, London.
- Burgman MA 1987 An analysis of the distribution of plants on granite outcrops in southern Western Australia using Mantel tests. Vegetatio 71:79-86.
- Bussell JD & James SH 1997 Rocks as museums of evolutionary processes. Journal of the Royal Society of Western Australia 80:221-229.
- Diels L 1906 Die Pflanzenwelt von West-Australien südlich des Wendekreises. Englemann, Leipzig.
- Dixon KW 1994 Towards integrated conservation of Australian endangered plants - the Western Australian model. Biodiversity and Conservation 3:148-159.
- Erickson R 1969 The Drummonds of Hawthornden. Lamb Paterson, Perth.
- Erickson R, George AS, Marchant NG & Morcombe MK 1973 Flowers and Plants of Western Australia. Reed Books, Chatswood, NSW.
- Gaff DF 1981 The biology of resurrection plants. In: The Biology of Australian Plants (eds JS Pate & AJ McComb). The University of Western Australia Press, Nedlands, 114-146.
- Grant V 1981 Plant Speciation. Colombia University Press, New York.
- Green JW 1985 Census of the Vascular Plants of Western Australia. Western Australian Herbarium, Department of Agriculture, Perth.
- Hopper SD 1979 Biogeographical aspects of speciation in the south west Australian flora. Annual Review of Ecology and Systematics 10:399-422.
- Hopper SD 1981 Honeyeaters and their winter food plants on granite rocks in the central wheatbelt of Western Australia. Australian Wildlife Research 8:187-97.
- Hopper SD 1992 Patterns of diversity at the population and species levels in south-west Australian mediterranean ecosystems. In: Biodiversity of Mediterranean Ecosystems in Australia (ed RJ Hobbs). Surrey Beatty & Sons, Sydney, 27-46.
- Hopper SD 1995 Evolutionary networks: natural hybridization and its conservation significance. In: Nature Conservation 4: the Role of Networks (eds DA Saunders, JL Craig & EM Mattiske). Surrey Beatty & Sons, Chipping Norton, NSW, 51-66.
- Hopper SD 1997 An Australian perspective on plant conservation biology in practice. In: Conservation Biology for the Coming Decade (eds PL Fiedler & P Kareiva). Chapman & Hall, New York. In press.
- Hopper SD & Burgman MA 1983 Cladistic and phenetic analyses of phylogenetic relationships among populations of *Eucalyptus caesia*. Australian Journal of Botany 31:161-172.
- Hopper SD, Harvey MS, Chappill JA, Main AR & Main BY 1996 The Western Australian biota as Gondwanan Heritage - a review. In: Gondwanan Heritage: Past, Present and Future of the Western Australian Biota (eds SD Hopper, JA Chappill, MS Harvey & AS George). Surrey Beatty & Sons, Chipping Norton, NSW, 1-46.
- Hopper SD, van Leeuwen S, Brown AP & Patrick SJ 1990 Western Australia's Endangered Flora. Department of Conservation and Land Management, Perth.
- Houle G 1990 Species-area relationship during primary succession in granite outcrop plant communities. American Journal of Botany 77:1433-1439.
- James SH 1965 Complex hybridity in *Isotoma petraea* I. The occurrence of interchange heterozygosity, autogamy and a balanced lethal system. Heredity 20:341-53.
- Keighery GJ 1995 How many weeds? In: Invasive Weeds and Regenerating Ecosystems in Western Australia (ed G Burke). Institute of Science & Technology Policy, Murdoch University, Perth, 8-12.
- Maiden JH 1909 Records of Western Australian Botanists. Journal of the Western Australian Natural History Society 2:5-26.
- Main BY 1967 Between Wodjil and Tor. Jacaranda Press, Brisbane, and Landfall Press, Perth.
- Main BY 1997 Granite outcrops: A collective ecosystem. Journal of the Royal Society of Western Australia 80:113-122.
- Marchant NG 1973 Species diversity in the south-western flora. Journal of the Royal Society of Western Australia 56:23-30.
- Mares MA 1997 The geobiological interface: Granitic outcrops as a selective force in mammalian evolution. Journal of the Royal Society of Western Australia 80:131-139.
- Moran GF & Hopper SD 1983 Genetic diversity and the insular population structure of the rare granite rock species, *Eucalyptus caesia* Benth. Australian Journal of Botany 31:161-172.
- Myers JS 1997 Geology of granites. Journal of the Royal Society of Western Australia 80:87-100.
- Newbey KR & Hnatiuk RJ 1985 Vegetation and flora. In: The Biological Survey of the Eastern Goldfields. Part 3. Jackson-Kalgoorlie Study Area. Records of the Western Australian Museum Supplement 23:11-38.
- Newbey KR, Keighery GJ & Hall NJ 1995 Vegetation and flora. In: The Biological Survey of the Eastern Goldfields. Part 11. Boorabbin-Southern Cross Study Area. Records of the Western Australian Museum Supplement 49:17-30.
- Ohlemüller R 1997 Biodiversity patterns of plant communities in shallow depressions on Western Australian granite outcrops (inselbergs). Diplomarbeit zur Erlangung des Grades eines Diplom-Biologen, University of Bonn.
- Ornduff R 1986 Comparative fecundity and population composition of heterostylous and non-heterostylous species of *Villarsia* (Menyanthaceae) in Western Australia. American Journal of Botany 73:282-286.
- Ornduff R 1987 Islands on islands: plant life on the granite outcrops of Western Australia. Harold L. Lyon Arboretum Lecture Number Fifteen. University of Hawaii Press, Honolulu.
- Ornduff R 1996 An unusual floral monomorphism in *Villarsia* (Menyanthaceae) and its proposed origin from distyly. In: Gondwanan Heritage: Past, Present and Future of the Western Australian Biota (eds SD Hopper, JA Chappill, MS Harvey & AS George). Surrey Beatty & Sons, Chipping Norton, NSW, 212-222.
- Pate JS & Dixon KW 1982 Tuberous, Cormous and Bulbous Plants. The University of Western Australia Press, Nedlands.
- Piggott PJ & Sage LW 1997 Remnant vegetation, priority flora and weed invasions at Yilliminning Rock. Journal of the Royal Society of Western Australia 80:201-208.
- Pignatti E & Pignatti S 1994 *Centrolepidi-Hydrocotyletea alatae*, a new class of ephemeral communities in Western Australia. Journal of Vegetation Science 5:55-62.

- Porembski S, Brown G & Barthlott W 1995 An inverted latitudinal gradient of plant diversity in shallow depressions on Ivorian inselbergs. *Vegetatio* 117:151-163.
- Porembski S, Seine R & Barthlott W 1997 Inselberg vegetation and the biodiversity of granite outcrops. *Journal of the Royal Society of Western Australia* 80:193-199.
- Sampson JF, Hopper SD & James SH 1988 Genetic diversity and the conservation of *Eucalyptus crucis* Maiden. *Australian Journal of Botany* 36:447-460.
- Schweinfurth U 1978 Beobachtungen an exponierten Standorten der sudwestaustralischen Küste. *Bot. Jahrb. Syst.* 99:168-187.
- Smith GG 1962 The flora of granite rocks of the Porongurup Range, South Western Australia. *Journal of the Royal Society of Western Australia* 45:18-23.
- Stace HM 1995 Protogyny, self-incompatibility and pollination in *Anthocercis gracilis* (Solanaceae). *Australian Journal of Botany* 43:451-459.
- Stoneburner A, Wyatt R, Catcheside DG & Stone IG 1993 Census of the mosses of Western Australia. *The Bryologist* 96:86-101.
- Stoneburner A & Wyatt R 1996 Ecological and phytogeographical characteristics of the mosses of Western Australia. In: *Gondwanan Heritage: Past, Present and Future of the Western Australian Biota* (eds SD Hopper, JA Chappill, MS Harvey & AS George). Surrey Beatty & Sons, Chipping Norton, NSW, 109-119.
- Walters TW & Wyatt R 1982 The vascular flora of granite outcrops in the Central Mineral Region of Texas. *Bulletin of the Torrey Botanical Club* 109:344-364.
- Wardell Johnson G & Williams M 1996 A floristic survey of the Tingle Mosaic, south-western Australia: applications in land use planning and management. *Journal of the Royal Society of Western Australia* 79:249-276.
- Wyatt R 1997 Reproductive ecology of granite outcrop plants from the south-eastern United States. *Journal of the Royal Society of Western Australia* 80:123-129.

Appendix 1

Alphabetical list of Western Australian plants (vascular and cryptogams combined) represented by specimens in the Western Australian Herbarium and for which habitat descriptions on labels included the words "granite outcrop".

List is current to 28 May 1996, from the WAHERB database of 413 000 specimens collected from all habitats throughout W A. Nomenclature follows Green (1985) and subsequent amendments included in the Western Australian Herbarium's WACENSUS database. Taxa are listed alphabetically by generic, specific and infraspecific names, with entry concluded by family name. Numbers are those assigned to vascular plant families as in Green (1985). Taxa who's family names are preceded by a letter rather than a number are cryptogams as follows: B - moss (bryophyte); H - liverwort (hepatic); L - lichen; F - macrofungus. Uncertainty as to taxonomic delimitation is given by *aff*, *sp*, ?, "unsorted", *sens. lat.*, or phrase name (e.g. *Form 'n'*, or "*walyahmoningenensis*". Well-delimited but unpublished taxa are identified by *ms* (for manuscript name). Full authorship of all names, published and unpublished, is given in Green (1985) and subsequent amendments included in WACENSUS.

Acacia aciphylla 163 Mimosaceae	Acacia jennerae 163 Mimosaceae
Acacia acuaria 163 Mimosaceae	Acacia jilberdingensis 163 Mimosaceae
Acacia acuminata subsp acuminata ms 163 Mimosaceae	Acacia lasiocalyx 163 Mimosaceae
Acacia acutata 163 Mimosaceae	Acacia leptopetala 163 Mimosaceae
Acacia aff cyperophylla 163 Mimosaceae	Acacia leptostachya 163 Mimosaceae
Acacia aff rhodophloia 163 Mimosaceae	Acacia ligulata 163 Mimosaceae
Acacia andrewsii 163 Mimosaceae	Acacia linophylla 163 Mimosaceae
Acacia aneura var aneura 163 Mimosaceae	Acacia littorea 163 Mimosaceae
Acacia aphylla 163 Mimosaceae	Acacia lysiphloia 163 Mimosaceae
Acacia appianata 163 Mimosaceae	Acacia merinthophora 163 Mimosaceae
Acacia assimilis subsp assimilis 163 Mimosaceae	Acacia merrallii 163 Mimosaceae
Acacia ayersiana var latifolia 163 Mimosaceae	Acacia microbotrys 163 Mimosaceae
Acacia browniana var browniana (typical variant) 163 Mimosaceae	Acacia monticola 163 Mimosaceae
Acacia browniana var obscura 163 Mimosaceae	Acacia multisiliqua 163 Mimosaceae
Acacia cochlearis 163 Mimosaceae	Acacia myrtifolia 163 Mimosaceae
Acacia conniana 163 Mimosaceae	Acacia olgana 163 Mimosaceae
Acacia constablei 163 Mimosaceae	Acacia prainii 163 Mimosaceae
Acacia coolgardiensis subsp coolgardiensis 163 Mimosaceae	Acacia pravifolia 163 Mimosaceae
Acacia coolgardiensis subsp effusa (Pedunculate variant) 163 Mimosaceae	Acacia pulchella subsp pulchella 163 Mimosaceae
Acacia costiniana 163 Mimosaceae	Acacia pulchella var goadbyi 163 Mimosaceae
Acacia cowaniana 163 Mimosaceae	Acacia punctulata ms 163 Mimosaceae
Acacia cracentis ms 163 Mimosaceae	Acacia ramulosa 163 Mimosaceae
Acacia crenulata ms 163 Mimosaceae	Acacia redolens 163 Mimosaceae
Acacia cuneifolia ms 163 Mimosaceae	Acacia restacea 163 Mimosaceae
Acacia cyclops 163 Mimosaceae	Acacia rigens 163 Mimosaceae
Acacia denticulosa 163 Mimosaceae	Acacia saligna 163 Mimosaceae
Acacia drummondii subsp elegans Porongurup variant (RJ Cumm ms) 163 Mimosaceae	Acacia scleroclada 163 Mimosaceae
Acacia ephedroides 163 Mimosaceae	Acacia sessilispica 163 Mimosaceae
Acacia ericifolia 163 Mimosaceae	Acacia sp 163 Mimosaceae
Acacia extensa 163 Mimosaceae	Acacia sp P58 (BR Maslin 3625) 163 Mimosaceae
Acacia fauntleroyi 163 Mimosaceae	Acacia sp P94 (JS Beard 6178) 163 Mimosaceae
Acacia flavipila var flavipila 163 Mimosaceae	Acacia stenoptera 163 Mimosaceae
Acacia granitcola ms 163 Mimosaceae	Acacia stowardii 163 Mimosaceae
Acacia hastulata 163 Mimosaceae	Acacia subcaerulea 163 Mimosaceae
Acacia hemiteles 163 Mimosaceae	Acacia subflexuosa subsp subflexuosa 163 Mimosaceae
Acacia heteroclita subsp heteroclita ms 163 Mimosaceae	Acacia tarculensis 163 Mimosaceae
Acacia heteroclita subsp valida ms 163 Mimosaceae	Acacia tetragonophylla 163 Mimosaceae
Acacia huegelii 163 Mimosaceae	Acacia trachycarpa 163 Mimosaceae

Acacia <i>tratmaniana</i>	163 Mimosaceae	Baeckea <i>crispiflora</i> ms 273 Myrtaceae
Acacia <i>trigonophylla</i>	163 Mimosaceae	Baeckea <i>maidenii</i> 273 Myrtaceae
Acacia <i>triplycha</i>	163 Mimosaceae	Baeckea <i>margarethae</i> 273 Myrtaceae
Acacia <i>urophylla</i>	163 Mimosaceae	Baeckea <i>margarethae</i> ms 273 Myrtaceae
Acacia <i>verrucula</i>	163 Mimosaceae	Baeckea <i>polyandra</i> 273 Myrtaceae
Acacia <i>viscidifolia</i>	163 Mimosaceae	Baeckea <i>recurva</i> ms 273 Myrtaceae
Acacia <i>willdenowiana</i>	163 Mimosaceae	Baeckea sp <i>Paynes Find</i> (S Patrick 1095) 273 Myrtaceae
Acacia ? <i>glaucissima</i>	163 Mimosaceae	Baeckea <i>tenuiramea</i> 273 Myrtaceae
Acacia ? <i>quadrimarginata</i>	163 Mimosaceae	Baeckea <i>tetragona</i> 273 Myrtaceae
Acacia Sect. Pluri. (microneurae, terete)	163 Mimosaceae	Banksia <i>seminuda</i> subsp <i>remanans</i> 090 Proteaceae
Acarospora <i>citrina</i> L	Acarosporaceae	Banksia <i>verticillata</i> 090 Proteaceae
Acaulon <i>integrifolium</i> B	Pottiaceae	Barbula <i>calycina</i> B Pottiaceae
Actinobole <i>oldfieldiana</i>	345 Asteraceae	Barbula <i>crinata</i> B Pottiaceae
Actinotus <i>leucocephalus</i>	281 Apiaceae	Bartsia <i>trixago</i> 316 Scrophulariaceae
Adenanthes <i>obovatus</i>	090 Proteaceae	Beaufortia <i>empetrifolia</i> 273 Myrtaceae
Agaricus sp F	Agaricaceae	Beaufortia <i>incana</i> 273 Myrtaceae
Agonis <i>hypericifolia</i>	273 Myrtaceae	Beaufortia <i>schaueri</i> 273 Myrtaceae
Agonis <i>linearifolia</i>	273 Myrtaceae	Blennospora <i>drummondii</i> 345 Asteraceae
Agonis <i>marginalis</i>	273 Myrtaceae	Boronia <i>albiflora</i> 175 Rutaceae
Agonis <i>parviceps</i>	273 Myrtaceae	Boronia <i>algida</i> 175 Rutaceae
Agrostocrinum <i>scabrum</i>	054F Anthericaceae	Boronia <i>busselliana</i> 175 Rutaceae
Aira <i>cupaniana</i>	031 Poaceae	Boronia <i>coerulescens</i> subsp "unsorted" 175 Rutaceae
Allocasuarina <i>campesiris</i>	070 Casuarinaceae	Boronia <i>crassifolia</i> 175 Rutaceae
Allocasuarina <i>decussata</i>	070 Casuarinaceae	Boronia <i>crenulata</i> var "unsorted" 175 Rutaceae
Allocasuarina <i>erioclathrys</i>	subsp <i>grossa</i> 070 Casuarinaceae	Boronia <i>crenulata</i> var <i>crenulata</i> 175 Rutaceae
Allocasuarina <i>huegeliana</i>	070 Casuarinaceae	Boronia <i>defoliatia</i> 175 Rutaceae
Allocasuarina <i>humilis</i>	070 Casuarinaceae	Boronia <i>gracilipes</i> 175 Rutaceae
Allocasuarina <i>thyoides</i>	070 Casuarinaceae	Boronia <i>inornata</i> subsp <i>inornata</i> 175 Rutaceae
Allocasuarina <i>trichodon</i>	070 Casuarinaceae	Boronia sp 175 Rutaceae
Alyogyne <i>hakeifolia</i>	221 Malvaceae	Boronia <i>subsessilis</i> 175 Rutaceae
Alyogyne <i>huegelii</i> var <i>huegelii</i>	ms 221 Malvaceae	Boronia <i>tetrandra</i> 175 Rutaceae
Alyogyne <i>huegelii</i> var <i>wrayae</i>	ms 221 Malvaceae	Borya <i>constricta</i> 054F Anthericaceae
Alyogyne <i>pinoniana</i>	221 Malvaceae	Borya <i>constricta</i> F Anthericaceae
Amanita sp F	Amanitaceae	Borya <i>laciniata</i> 054F Anthericaceae
Amaranthus <i>mitchellii</i>	106 Amaranthaceae	Borya <i>longiscapa</i> 054F Anthericaceae
Amaranthus <i>pallidiflorus</i>	106 Amaranthaceae	Borya <i>nitida</i> 054F Anthericaceae
Amaranthus sp (Silent Grove)	RJC 6532 106 Amaranthaceae	Borya <i>scirpoidea</i> 054F Anthericaceae
Amperea <i>simulans</i>	185 Euphorbiaceae	Borya <i>sphaerocephala</i> 054F Anthericaceae
Amphipogon <i>caricinus</i>	031 Poaceae	Borya <i>sphaerocephala</i> sens lat 054F Anthericaceae
Amyema <i>gibberula</i> var <i>tatei</i>	097 Loranthaceae	Bossiaea <i>dentata</i> 163 Papilionaceae
Anagallis <i>arvensis</i>	293 Primulaceae	Bossiaea <i>eriocarpa</i> 165 Papilionaceae
Anarthria <i>prolifera</i>	039 Restionaceae	Bossiaea <i>linophylla</i> 165 Papilionaceae
Andersonia aff <i>auriculata</i>	288 Epacridaceae	Brachychiton <i>grandiflorus</i> 223 Sterculiaceae
Andersonia aff <i>lehmanniana</i>	288 Epacridaceae	Brachychiton <i>gregorii</i> 223 Sterculiaceae
Andersonia <i>aristata</i>	288 Epacridaceae	Brachynemurum <i>preissianum</i> B Bryaceae
Andersonia <i>lehmanniana</i>	288 Epacridaceae	Brachyscome <i>ciliaris</i> 345 Asteraceae
Andersonia <i>parvifolia</i>	288 Epacridaceae	Brachyscome <i>iberidifolia</i> 345 Asteraceae
Andersonia sp	288 Epacridaceae	Brachyscome <i>perpusilla</i> 345 Asteraceae
Andersonia <i>sprengelioidea</i>	288 Epacridaceae	Brachyscome sp 345 Asteraceae
Angianthus <i>tomentosus</i>	345 Asteraceae	Breutelia <i>affinis</i> B Bartramiaceae
Anthocercis <i>anisantha</i>	subsp <i>anisantha</i> 315 Solanaceae	Briza <i>maxima</i> 031 Poaceae
Anthocercis <i>viscosa</i>	subsp <i>viscosa</i> 315 Solanaceae	Briza <i>minor</i> 031 Poaceae
Anthurus sp F	Clathraceae	Bromus <i>rubens</i> 031 Poaceae
Aphelia <i>brizula</i>	040 Centrolepidaceae	Bruchia <i>brevipes</i> B Dicranaceae
Aphelia <i>cyperoides</i>	040 Centrolepidaceae	Bryum <i>argenteum</i> B Bryaceae
Aphelia <i>nutans</i>	040 Centrolepidaceae	Bryum <i>billardieri</i> B Bryaceae
Aristida <i>contorta</i>	031 Poaceae	Bryum <i>caespiticium</i> B Bryaceae
Arthropodium <i>curvipes</i>	054F Anthericaceae	Bryum <i>campylothecium</i> B Bryaceae
Arthropodium <i>dyeri</i>	054F Anthericaceae	Bryum <i>capillare</i> B Bryaceae
Asplenium <i>aethiopicum</i>	011E Aspleniaceae	Bryum <i>cheelii</i> B Bryaceae
Asplenium <i>flabellifolium</i>	011E Aspleniaceae	Bryum <i>chrysoneuron</i> B Bryaceae
Astarea <i>ambigua</i>	273 Myrtaceae	Bryum <i>dichotomum</i> B Bryaceae
Astarea <i>fascicularis</i>	273 Myrtaceae	Bryum <i>pachytheca</i> B Bryaceae
Astarea sp Mt Johnston (AR Annals 5645)	273 Myrtaceae	Bryum sp B <i>Catcheside</i> B Bryaceae
Asterella <i>drummondii</i>	H Hepaticae	Bryum <i>torquescens</i> B Bryaceae
Asterolasia <i>pallida</i>	subsp "unsorted" 175 Rutaceae	Bulbine <i>semibarbata</i> 054G Asphodelaceae
Astroloma <i>ciliatum</i>	288 Epacridaceae	Burchardia <i>multiflora</i> 054J Colchicaceae
Astroloma <i>epacridis</i>	288 Epacridaceae	Bursaria <i>occidentalis</i> 152 Pittosporaceae
Astroloma <i>pallidum</i>	288 Epacridaceae	Caesia <i>micrantha</i> 054F Anthericaceae
Astroloma <i>prostratum</i>	288 Epacridaceae	Caladenia <i>attingen</i> subsp <i>attingen</i> ms 066 Orchidaceae
Astroloma <i>serratifolium</i>	288 Epacridaceae	Caladenia <i>attingen</i> subsp <i>gracillima</i> ms 066 Orchidaceae
Astroloma sp	288 Epacridaceae	Caladenia <i>brevisura</i> ms 066 Orchidaceae
Baeckea "walyahmoningensis"	273 Myrtaceae	Caladenia <i>brownii</i> ms 066 Orchidaceae
Baeckea <i>behrii</i>	273 Myrtaceae	Caladenia <i>caesarea</i> subsp <i>maritima</i> ms 066 Orchidaceae
Baeckea <i>camphorosmae</i>	273 Myrtaceae	Caladenia <i>caesarea</i> subsp <i>transiens</i> ms 066 Orchidaceae
Baeckea <i>crispiflora</i>	273 Myrtaceae	Caladenia <i>citrina</i> ms 066 Orchidaceae

<i>Caladenia doutchiae</i> 066 Orchidaceae	<i>Centaurium erythraea</i> 303 Gentianaceae
<i>Caladenia extans</i> ms 066 Orchidaceae	<i>Centaurium</i> sp 303 Gentianaceae
<i>Caladenia falcata</i> 066 Orchidaceae	<i>Centranthus ruber</i> 334 Valerianaceae
<i>Caladenia filifera</i> 066 Orchidaceae	<i>Centrolepis alepyrooides</i> 040 Centrolepidaceae
<i>Caladenia flaccida</i> subsp <i>flaccida</i> ms 066 Orchidaceae	<i>Centrolepis aristata</i> 040 Centrolepidaceae
<i>Caladenia flaccida</i> subsp <i>pulchra</i> ms 066 Orchidaceae	<i>Centrolepis drummondiana</i> 040 Centrolepidaceae
<i>Caladenia flava</i> subsp <i>flava</i> ms 066 Orchidaceae	<i>Centrolepis eremica</i> 040 Centrolepidaceae
<i>Caladenia flava</i> subsp <i>maculata</i> ms 066 Orchidaceae	<i>Centrolepis glabra</i> 040 Centrolepidaceae
<i>Caladenia flava</i> subsp <i>sylvestris</i> ms 066 Orchidaceae	<i>Centrolepis pilosa</i> 040 Centrolepidaceae
<i>Caladenia footiana</i> ms 066 Orchidaceae	<i>Centrolepis polygyna</i> 040 Centrolepidaceae
<i>Caladenia granitiora</i> ms 066 Orchidaceae	<i>Centrolepis strigosa</i> subsp <i>rupestris</i> 040 Centrolepidaceae
<i>Caladenia heberleana</i> ms 066 Orchidaceae	<i>Centrolepis strigosa</i> subsp <i>rupestris</i> ms 040 Centrolepidaceae
<i>Caladenia hirta</i> subsp <i>rosea</i> ms 066 Orchidaceae	<i>Centrolepis strigosa</i> subsp <i>strigosa</i> 040 Centrolepidaceae
<i>Caladenia hoffmannii</i> subsp <i>graniticola</i> ms 066 Orchidaceae	<i>Cephalozia arctica</i> subsp <i>subantarctica</i> H Cephaloziellaceae
<i>Caladenia incensa</i> ms 066 Orchidaceae	<i>Chamaescilla corymbosa</i> var <i>corymbosa</i> 054F Anthericaceae
<i>Caladenia incrassata</i> ms 066 Orchidaceae	<i>Chamaescilla corymbosa</i> var <i>latifolia</i> 054F Anthericaceae
<i>Caladenia infundibularis</i> 066 Orchidaceae	<i>Chamelaicum ciliatum</i> 273 Myrtaceae
<i>Caladenia integra</i> 066 Orchidaceae	<i>Chamelaicum floriferum</i> ms 273 Myrtaceae
<i>Caladenia latifolia</i> 066 Orchidaceae	<i>Chamelaicum floriferum</i> subsp <i>diffusum</i> ms 273 Myrtaceae
<i>Caladenia lobata</i> 066 Orchidaceae	<i>Chamelaicum floriferum</i> subsp <i>floriferum</i> ms 273 Myrtaceae
<i>Caladenia longicauda</i> subsp <i>clivicola</i> ms 066 Orchidaceae	<i>Chamelaicum forrestii</i> 273 Myrtaceae
<i>Caladenia longicauda</i> subsp <i>eminens</i> ms 066 Orchidaceae	<i>Chamelaicum forrestii</i> subsp <i>forrestii</i> ms 273 Myrtaceae
<i>Caladenia longicauda</i> subsp <i>longicauda</i> ms 066 Orchidaceae	<i>Chamelaicum</i> sp <i>Yalgoo</i> (Y Chadwick 1816) ms 273 Myrtaceae
<i>Caladenia longicauda</i> subsp <i>rigida</i> ms 066 Orchidaceae	<i>Chamomilla</i> sp F Hymenogastraceae
<i>Caladenia longilabata</i> 066 Orchidaceae	<i>Cheilanthes</i> aff <i>austrotenuifolia</i> 007 Adiantaceae
<i>Caladenia macrostylis</i> 066 Orchidaceae	<i>Cheilanthes austrotenuifolia</i> 007 Adiantaceae
<i>Caladenia marginata</i> 066 Orchidaceae	<i>Cheilanthes distans</i> 007 Adiantaceae
<i>Caladenia microchila</i> ms 066 Orchidaceae	<i>Cheilanthes lasiophylla</i> 007 Adiantaceae
<i>Caladenia multiclavia</i> 066 Orchidaceae	<i>Cheilanthes sieberi</i> subsp <i>sieberi</i> 007 Adiantaceae
<i>Caladenia nivalis</i> ms 066 Orchidaceae	<i>Cheiranthera filifolia</i> var <i>filifolia</i> 152 Pittosporaceae
<i>Caladenia pachychila</i> ms 066 Orchidaceae	<i>Chiloscyphus semiteres</i> H Geocalycaceae
<i>Caladenia pendens</i> ms 066 Orchidaceae	<i>Chloris truncata</i> 031 Poaceae
<i>Caladenia pholcoidea</i> ms 066 Orchidaceae	<i>Chorizandra enodis</i> 032 Cyperaceae
<i>Caladenia polychroma</i> ms 066 Orchidaceae	<i>Chorizema aciculare</i> subsp <i>aciculare</i> 165 Papilionaceae
<i>Caladenia radialis</i> 066 Orchidaceae	<i>Chorizema cordatum</i> 165 Papilionaceae
<i>Caladenia reptans</i> subsp <i>impensa</i> ms 066 Orchidaceae	<i>Chorizema dicksonii</i> 165 Papilionaceae
<i>Caladenia reptans</i> subsp <i>reptans</i> ms 066 Orchidaceae	<i>Chorizema nervosum</i> 165 Papilionaceae
<i>Caladenia rhomboidiformis</i> 066 Orchidaceae	<i>Chorizema retrorsum</i> 165 Papilionaceae
<i>Caladenia roei</i> 066 Orchidaceae	<i>Chrysoccephalum apiculatum</i> 345 Asteraceae
<i>Caladenia saccharata</i> 066 Orchidaceae	<i>Chrysoccephalum ramosissimum</i> 345 Asteraceae
<i>Caladenia serotina</i> ms 066 Orchidaceae	<i>Chrysoccephalum semicalvum</i> 345 Asteraceae
<i>Caladenia sigmae</i> 066 Orchidaceae	<i>Chthonocephalus pseudexav</i> 345 Asteraceae
<i>Caladenia splendens</i> ms 066 Orchidaceae	<i>Cladonia aggregata</i> L Cladiaceae
<i>Caladenia hiemalis</i> ms 066 Orchidaceae	<i>Cladonia ferdinandii</i> L Cladiaceae
<i>Caladenia horistes</i> ms 066 Orchidaceae	<i>Cladonia sullivani</i> L Cladiaceae
<i>Caladenia pendens</i> subsp <i>pendens</i> ms 066 Orchidaceae	<i>Cladonia capitellata</i> L Cladoniaceae
<i>Caladenia remota</i> subsp <i>remota</i> ms 066 Orchidaceae	<i>Clematicissus angustissima</i> 216 Vitaceae
<i>Caladenia pendens</i> subsp <i>talbotii</i> ms 066 Orchidaceae	<i>Clematis pubescens</i> 119 Ranunculaceae
<i>Caladenia voigtii</i> ms 066 Orchidaceae	<i>Cleome tetrandra</i> var <i>tetrandra</i> 137A Capparaceae
<i>Calandrinia calyptrata</i> 111 Portulacaceae	<i>Cleome uncifera</i> 137A Capparaceae
<i>Calandrinia ptychosperma</i> 111 Portulacaceae	<i>Coccocarpia erythroxili</i> L Coccocarpiaceae
<i>Calectasia grandiflora</i> 054C Dasypogonaceae	<i>Cochlospermum gillivrai</i> 241 Cochlospermaceae
<i>Callitricha stagnalis</i> 186 Callitrichaceae	<i>Comesperma calymega</i> 183 Polygalaceae
<i>Callitris glauccophylla</i> 018 Cupressaceae	<i>Comesperma confertum</i> 183 Polygalaceae
<i>Callitris preissii</i> subsp "unsorted" 018 Cupressaceae	<i>Comesperma integrerrimum</i> 183 Polygalaceae
<i>Calocephalus multiflorus</i> 345 Asteraceae	<i>Comesperma spinosum</i> 183 Polygalaceae
<i>Caloplaca</i> sp L Teloschistaceae	<i>Comesperma volubile</i> 183 Polygalaceae
<i>Calothamnus quadrifidus</i> var "unsorted" 273 Myrtaceae	<i>Commersonia crispa</i> 223 Sterculiaceae
<i>Calothamnus quadrifidus</i> var <i>Esperance District</i> (AE Orchard 1676) 273 Myrtaceae	<i>Conospermum caeruleum</i> 090 Proteaceae
<i>Calothamnus rupestris</i> 273 Myrtaceae	<i>Conostylis aculeata</i> subsp <i>aculeata</i> 055 Haemodoraceae
<i>Calothamnus schaueri</i> 273 Myrtaceae	<i>Conostylis bealiana</i> 055 Haemodoraceae
<i>Calothamnus</i> sp 273 Myrtaceae	<i>Conostylis prolifera</i> 055 Haemodoraceae
<i>Calothamnus tuberosus</i> 273 Myrtaceae	<i>Conostylis setigera</i> 055 Haemodoraceae
<i>Calothamnus villosus</i> 273 Myrtaceae	<i>Conostylis setigera</i> subsp <i>setigera</i> 055 Haemodoraceae
<i>Calycopeplus paucifolius</i> 185 Euphorbiaceae	<i>Conostylis setosa</i> 055 Haemodoraceae
<i>Calytrix acutifolia</i> 273 Myrtaceae	<i>Corchorus elachocarpus</i> 220 Tiliaceae
<i>Calytrix angulata</i> 273 Myrtaceae	<i>Corybas recurvus</i> 066 Orchidaceae
<i>Calytrix decandra</i> 273 Myrtaceae	<i>Cotula coronopifolia</i> 345 Asteraceae
<i>Calytrix depressa</i> 273 Myrtaceae	<i>Cotula cotuloides</i> 345 Asteraceae
<i>Calytrix fraseri</i> 273 Myrtaceae	<i>Craspedia variabilis</i> 345 Asteraceae
<i>Calytrix glutinosa</i> 273 Myrtaceae	<i>Crassula colorata</i> var <i>acuminata</i> 149 Crassulaceae
<i>Calytrix leschenaultii</i> 273 Myrtaceae	<i>Crassula decumbens</i> var <i>decumbens</i> 149 Crassulaceae
<i>Calytrix tetragona</i> 273 Myrtaceae	<i>Crassula exserta</i> 149 Crassulaceae
<i>Campylopus australis</i> B Dicranaceae	<i>Crassula natans</i> var <i>minus</i> 149 Crassulaceae
<i>Campylopus bicolor</i> B Dicranaceae	<i>Crassula pedicellosa</i> 149 Crassulaceae
<i>Campylopus flindersii</i> B Dicranaceae	<i>Crassula peduncularis</i> 149 Crassulaceae
<i>Campylopus introflexus</i> B Dicranaceae	<i>Crassula sieberiana</i> subsp <i>tetramera</i> 149 Crassulaceae
<i>Carpobrotus aequilaterus</i> 110 Aizoaceae	<i>Crassula</i> sp 149 Crassulaceae
<i>Cassytha glabella</i> forma <i>dispar</i> 131 Lauraceae	<i>Crossidium davidai</i> B Pottiaceae
<i>Cassytha racemosa</i> 131 Lauraceae	<i>Crotalaria</i> sp 165 Papilionaceae

<i>Cryptandra arbutilflora</i> var. <i>arbutilflora</i> 215 Rhamnaceae	<i>Dodonaea viscosa</i> subsp. <i>angustissima</i> 207 Sapindaceae
<i>Cryptandra congesta</i> 215 Rhamnaceae	<i>Dodonaea viscosa</i> subsp. <i>mucronata</i> 207 Sapindaceae
<i>Cryptandra pungens</i> 215 Rhamnaceae	<i>Dodonaea viscosa</i> subsp. <i>spatulata/angustissima</i> 207 Sapindaceae
<i>Cryptandra wilsonii</i> 215 Rhamnaceae	<i>Dodonaea viscosa</i> subsp. <i>spatulata</i> 207 Sapindaceae
<i>Cryptostylis ovata</i> 066 Orchidaceae	<i>Drakonorchis barbrossa</i> ms 066 Orchidaceae
<i>Cuscuta epithymum</i> 307A Cuscutaceae	<i>Drakonorchis drakeoides</i> ms 066 Orchidaceae
<i>Cyanicula amplexans</i> ms 066 Orchidaceae	<i>Drakonorchis mesocera</i> ms 066 Orchidaceae
<i>Cyanicula ashbyae</i> ms 066 Orchidaceae	<i>Drosa andersoniana</i> 143 Droseraceae
<i>Cyanicula caerulea</i> subsp. <i>apertala</i> ms 066 Orchidaceae	<i>Drosa bulbosa</i> subsp. <i>bulbosa</i> 143 Droseraceae
<i>Cyanicula deformis</i> ms 066 Orchidaceae	<i>Drosa erythrogynae</i> 143 Droseraceae
<i>Cyanicula fragrans</i> ms 066 Orchidaceae	<i>Drosa gigantea</i> subsp. <i>gigantea</i> 143 Droseraceae
<i>Cyanicula gemmata</i> ms 066 Orchidaceae	<i>Drosa glanduligera</i> 143 Droseraceae
<i>Cyanicula sericea</i> ms 066 Orchidaceae	<i>Drosa heterophylla</i> 143 Droseraceae
<i>Cycas pruinosa</i> 016 Cycadaceae	<i>Drosa leucoblasta</i> 143 Droseraceae
<i>Cynoglossum australe</i> var. "unsorted" 310 Boraginaceae	<i>Drosa lowriei</i> 143 Droseraceae
<i>Cyperus centralis</i> 032 Cyperaceae	<i>Drosa macrantha</i> subsp. <i>macrantha</i> 143 Droseraceae
<i>Cyperus cunninghamii</i> subsp. <i>cunninghamii</i> 032 Cyperaceae	<i>Drosa menziesii</i> 143 Droseraceae
<i>Cyperus tenellus</i> 032 Cyperaceae	<i>Drosa menziesii</i> subsp. <i>menziesii</i> 143 Droseraceae
<i>Cyphanthera microphylla</i> 315 Solanaceae	<i>Drosa menziesii</i> subsp. <i>penicillaris</i> 143 Droseraceae
<i>Cyrtostylis huegelii</i> var. 066 Orchidaceae	<i>Drosa microphylla</i> 143 Droseraceae
<i>Cyrtostylis robusta</i> <i>reniformis</i> 066 Orchidaceae	<i>Drosa peltata</i> 143 Droseraceae
<i>Dampiera alata</i> 341 Goodeniaceae	<i>Drosa ramellosa</i> 143 Droseraceae
<i>Dampiera altissima</i> 341 Goodeniaceae	<i>Drosa</i> sp. 143 Droseraceae
<i>Dampiera fasciculata</i> 341 Goodeniaceae	<i>Drosa stolonifera</i> 143 Droseraceae
<i>Dampiera haematotrichia</i> subsp. <i>dura</i> 341 Goodeniaceae	<i>Drosa stolonifera</i> subsp. <i>rupicola</i> 143 Droseraceae
<i>Dampiera lavandulacea</i> 341 Goodeniaceae	<i>Drosa stolonifera</i> subsp. <i>stolonifera</i> 143 Droseraceae
<i>Dampiera linearis</i> 341 Goodeniaceae	<i>Drosa subhirtella</i> subsp. <i>moorei</i> 143 Droseraceae
<i>Dampiera</i> sp. 341 Goodeniaceae	<i>Drosa subhirtella</i> subsp. <i>subhirtella</i> 143 Droseraceae
<i>Dampiera stenostachya</i> 341 Goodeniaceae	<i>Dryandra armata</i> 090 Proteaceae
<i>Danthonia caespitosa</i> 031 Poaceae	<i>Dryandra armata</i> var. <i>armata</i> 090 Proteaceae
<i>Danthonia</i> sp. 031 Poaceae	<i>Dryandra cuneata</i> 090 Proteaceae
<i>Darwinia citriodora</i> 273 Myrtaceae	<i>Dryandra nivea</i> subsp. <i>nivea</i> ms 090 Proteaceae
<i>Darwinia diosmoides</i> 273 Myrtaceae	<i>Dysphania glomulifera</i> subsp. <i>eremaea</i> 105 Chenopodiaceae
<i>Darwinia thymoides</i> 273 Myrtaceae	<i>Eccremidium arcutatum</i> B Ditrichaceae
<i>Darwinia vestita</i> 273 Myrtaceae	<i>Eccremidium pulchellum</i> B Ditrichaceae
<i>Daucus glochidiatus</i> 281 Apiaceae	<i>Ecdeiocolea monostachya</i> 039 Restionaceae
<i>Daviesia benthamii</i> subsp. <i>acanthoclona</i> ms 165 Papilionaceae	<i>Elatine gratioloides</i> 235 Elatinaceae
<i>Daviesia croniiniana</i> 165 Papilionaceae	<i>Elythranthera brunonis</i> 066 Orchidaceae
<i>Daviesia decurrens</i> 165 Papilionaceae	<i>Elythranthera emarginata</i> 066 Orchidaceae
<i>Daviesia horrida</i> 165 Papilionaceae	<i>Enchytraea tomentosa</i> var. <i>tomentosa</i> 105 Chenopodiaceae
<i>Daviesia incrassata</i> subsp. <i>reversifolia</i> 165 Papilionaceae	<i>Endocarpion</i> sp. (<i>crassisporum vel macrosporum</i>) L Verrucariaceae
<i>Daviesia inflata</i> 165 Papilionaceae	<i>Epheba lanata</i> L Lichenaceae
<i>Desmatodon convolutus</i> B Pottiaceae	<i>Ephemerum cristatum</i> B Ephemeraceae
<i>Desmatodon recurvatus</i> B Pottiaceae	<i>Eremophila alternifolia</i> 326 Myoporaceae
<i>Desmatodon</i> sp. nov X B Pottiaceae	<i>Eremophila clarkei</i> 326 Myoporaceae
<i>Desmodcladus aspera</i> ms 039 Restionaceae	<i>Eremophila decipiens</i> subsp. "unsorted" 326 Myoporaceae
<i>Dichondra repens</i> 307 Convolvulaceae	<i>Eremophila decipiens</i> subsp. <i>decipiens</i> ms 326 Myoporaceae
<i>Dichopogon capillipes</i> 054F Anthericaceae	<i>Eremophila deserti</i> 326 Myoporaceae
<i>Dicranoloma diaphanoneurum</i> B Dicranaceae	<i>Eremophila exilifolia</i> 326 Myoporaceae
<i>Didymodon torquatus</i> B Pottiaceae	<i>Eremophila forrestii</i> subsp. "unsorted" 326 Myoporaceae
<i>Digitaria brownii</i> 031 Poaceae	<i>Eremophila glabra</i> subsp. "unsorted" 326 Myoporaceae
<i>Dillwynia</i> sp. A Perth Flora (R Coveny 8036) 165 Papilionaceae	<i>Eremophila glutinosa</i> 326 Myoporaceae
<i>Dioscorea hastifolia</i> 059 Dioscoreaceae	<i>Eremophila lehmanniana</i> 326 Myoporaceae
<i>Diplocyclos palmatus</i> 337 Cucurbitaceae	<i>Eremophila oldfieldii</i> subsp. <i>angustifolia</i> ms 326 Myoporaceae
<i>Diplolaena andrewsii</i> 175 Rutaceae	<i>Eremophila oppositifolia</i> subsp. <i>angustifolia</i> ms 326 Myoporaceae
<i>Diplolaena graniticola</i> ms 175 Rutaceae	<i>Eremophila pachyphylla</i> 326 Myoporaceae
<i>Diplolaena microcephala</i> 175 Rutaceae	<i>Eremophila phillipsii</i> 326 Myoporaceae
<i>Diplolaena velutina</i> ms 175 Rutaceae	<i>Eremophila platycalyx</i> subsp. <i>platycalyx</i> ms 326 Myoporaceae
<i>Diploschistes</i> sp. L Thelotremaeaceae	<i>Eremophila scoparia</i> 326 Myoporaceae
<i>Diuris</i> aff. <i>longifolia</i> 066 Orchidaceae	<i>Eremophila serrulata</i> 326 Myoporaceae
<i>Diuris brumalis</i> 066 Orchidaceae	<i>Eremophila</i> sp. 326 Myoporaceae
<i>Diuris conspicillata</i> 066 Orchidaceae	<i>Eriachne pulchella</i> subsp. <i>pulchella</i> 031 Poaceae
<i>Diuris laevis</i> 066 Orchidaceae	<i>Eriochilus dilatatus</i> subsp. <i>dilatatus</i> ms 066 Orchidaceae
<i>Diuris laxiflora</i> 066 Orchidaceae	<i>Eriochilus dilatatus</i> subsp. <i>multiflorus</i> ms 066 Orchidaceae
<i>Diuris longifolia</i> 066 Orchidaceae	<i>Eriochilus dilatatus</i> subsp. <i>undulatus</i> ms 066 Orchidaceae
<i>Diuris maculata</i> 066 Orchidaceae	<i>Eriochilus helonomus</i> ms 066 Orchidaceae
<i>Diuris picta</i> 066 Orchidaceae	<i>Eriochilus pulchellus</i> ms 066 Orchidaceae
<i>Diuris pulchella</i> 066 Orchidaceae	<i>Eriochilus scaber</i> 066 Orchidaceae
<i>Diuris recurva</i> 066 Orchidaceae	<i>Eriostemon brucei</i> subsp. <i>brucei</i> 175 Rutaceae
<i>Diuris setacea</i> 066 Orchidaceae	<i>Eriostemon linearis</i> 175 Rutaceae
<i>Dodonaea bursariifolia</i> 207 Sapindaceae	<i>Erodium cicutarium</i> 167 Geraniaceae
<i>Dodonaea caespitosa</i> 207 Sapindaceae	<i>Eucalyptus acies</i> 273 Myrtaceae
<i>Dodonaea ceratocarpa</i> 207 Sapindaceae	<i>Eucalyptus aff. lane-poolei</i> 273 Myrtaceae
<i>Dodonaea inaequifolia</i> 207 Sapindaceae	<i>Eucalyptus angulosa</i> 273 Myrtaceae
<i>Dodonaea lobulata</i> 207 Sapindaceae	<i>Eucalyptus arachnaea</i> 273 Myrtaceae
<i>Dodonaea microzyga</i> var. <i>acrolobata</i> 207 Sapindaceae	<i>Eucalyptus aspratilis</i> 273 Myrtaceae
<i>Dodonaea petiolaris</i> 207 Sapindaceae	<i>Eucalyptus brachyphylla</i> 273 Myrtaceae
<i>Dodonaea pinifolia</i> 207 Sapindaceae	<i>Eucalyptus brevipes</i> 273 Myrtaceae
<i>Dodonaea ptarmicaefolia</i> 207 Sapindaceae	<i>Eucalyptus caesia</i> subsp. <i>caesia</i> 273 Myrtaceae
<i>Dodonaea stenozyga</i> 207 Sapindaceae	<i>Eucalyptus calycogona</i> var. <i>calycogona</i> 273 Myrtaceae

<i>Eucalyptus cornuta</i> 273 Myrtaceae	<i>Gnephosis tenuissima</i> 345 Asteraceae
<i>Eucalyptus crucis</i> subsp <i>crucis</i> 273 Myrtaceae	<i>Gompholobium knightianum</i> 165 Papilionaceae
<i>Eucalyptus crucis</i> subsp <i>lanceolata</i> 273 Myrtaceae	<i>Gompholobium marginatum</i> 165 Papilionaceae
<i>Eucalyptus decipiens</i> 273 Myrtaceae	<i>Gompholobium polymorphum</i> 165 Papilionaceae
<i>Eucalyptus decipiens</i> subsp <i>adesmophloia</i> 273 Myrtaceae	<i>Gompholobium venustum</i> 165 Papilionaceae
<i>Eucalyptus decipiens</i> subsp <i>chalara</i> 273 Myrtaceae	<i>Gonocarpus nodulosus</i> 276 Haloragaceae
<i>Eucalyptus doratoxylon</i> 273 Myrtaceae	<i>Gonocarpus paniculatus</i> 276 Haloragaceae
<i>Eucalyptus drummondii</i> 273 Myrtaceae	<i>Gonocarpus scordioides</i> 276 Haloragaceae
<i>Eucalyptus eremophila</i> 273 Myrtaceae	<i>Gonocarpus</i> sp 276 Haloragaceae
<i>Eucalyptus eremophila</i> hybrid 273 Myrtaceae	<i>Goodenia affinis</i> 341 Goodeniaceae
<i>Eucalyptus flocktoniae</i> 273 Myrtaceae	<i>Goodenia caerulea</i> 341 Goodeniaceae
<i>Eucalyptus gratiae</i> 273 Myrtaceae	<i>Goodenia concinna</i> 341 Goodeniaceae
<i>Eucalyptus incrassata</i> 273 Myrtaceae	<i>Goodenia recursiva</i> 341 Goodeniaceae
<i>Eucalyptus insularis</i> 273 Myrtaceae	<i>Goodenia havilandii</i> 341 Goodeniaceae
<i>Eucalyptus kruseana</i> 273 Myrtaceae	<i>Goodenia helmsii</i> 341 Goodeniaceae
<i>Eucalyptus lehmannii</i> 273 Myrtaceae	<i>Goodenia incana</i> 341 Goodeniaceae
<i>Eucalyptus loxophleba</i> subsp <i>lissophloia</i> 273 Myrtaceae	<i>Goodenia micrantha</i> 341 Goodeniaceae
<i>Eucalyptus megacarpa</i> 273 Myrtaceae	<i>Goodenia muelleriana</i> 341 Goodeniaceae
<i>Eucalyptus melanoxylon</i> 273 Myrtaceae	<i>Goodenia pulchella</i> 341 Goodeniaceae
<i>Eucalyptus orbifolia</i> 273 Myrtaceae	<i>Goodenia quadrilocularis</i> 341 Goodeniaceae
<i>Eucalyptus patens</i> 273 Myrtaceae	<i>Goodenia scapigera</i> 341 Goodeniaceae
<i>Eucalyptus petraea</i> 273 Myrtaceae	<i>Goodenia stenophylla</i> 341 Goodeniaceae
<i>Eucalyptus phaenophylla</i> subsp <i>interjacens</i> 273 Myrtaceae	<i>Goodenia tripartita</i> 341 Goodeniaceae
<i>Eucalyptus pilleata</i> 273 Myrtaceae	<i>Granitites intangendus</i> 215 Rhhamnaceae
<i>Eucalyptus polycycla</i> 273 Myrtaceae	<i>Grevillea</i> aff <i>concinna</i> 090 Proteaceae
<i>Eucalyptus prava</i> ms 273 Myrtaceae	<i>Grevillea astericosa</i> 090 Proteaceae
<i>Eucalyptus subulata</i> 273 Myrtaceae	<i>Grevillea bipinnatifida</i> 090 Proteaceae
<i>Eucalyptus sheathiana</i> 273 Myrtaceae	<i>Grevillea centristigma</i> 090 Proteaceae
<i>Eucalyptus sporadica</i> subsp <i>sporadica</i> ms 273 Myrtaceae	<i>Grevillea christinae</i> 090 Proteaceae
<i>Eucalyptus suggrands</i> subsp <i>alipes</i> 273 Myrtaceae	<i>Grevillea cirsifolia</i> 090 Proteaceae
<i>Eucalyptus suggrands</i> subsp <i>biving</i> (PJW 588) 273 Myrtaceae	<i>Grevillea concinna</i> subsp <i>concinna</i> 090 Proteaceae
<i>Eucalyptus tephrodes</i> ms 273 Myrtaceae	<i>Grevillea concinna</i> subsp <i>leemanniana</i> 090 Proteaceae
<i>Eucalyptus tetraptera</i> 273 Myrtaceae	<i>Grevillea diversifolia</i> subsp <i>subtersericata</i> 090 Proteaceae
<i>Eucalyptus uncinata</i> 273 Myrtaceae	<i>Grevillea dolichopoda</i> 090 Proteaceae
<i>Eucalyptus virginiae</i> ms 273 Myrtaceae	<i>Grevillea fuscolutea</i> 090 Proteaceae
<i>Euphorbia coghlani</i> 185 Euphorbiaceae	<i>Grevillea granulosa</i> 090 Proteaceae
<i>Euphorbia</i> sp 185 Euphorbiaceae	<i>Grevillea huegelii</i> 090 Proteaceae
<i>Eutaxia microphylla</i> 165 Papilionaceae	<i>Grevillea leptobotrys</i> 090 Proteaceae
<i>Eutaxia obovata</i> 165 Papilionaceae	<i>Grevillea manglesii</i> subsp <i>manglesii</i> 090 Proteaceae
<i>Ficus platypoda</i> var <i>minor</i> 087 Moraceae	<i>Grevillea nana</i> subsp <i>nana</i> 090 Proteaceae
<i>Filago gallica</i> 345 Asteraceae	<i>Grevillea paniculata</i> (Form 'n') 090 Proteaceae
<i>Fimbristylis dichotoma</i> (desert form) 032 Cyperaceae	<i>Grevillea parallela</i> 090 Proteaceae
<i>Fimbristylis neilsonii</i> 032 Cyperaceae	<i>Grevillea pauciflora</i> subsp <i>psilophylla</i> 090 Proteaceae
<i>Fissidens asplenoides</i> B Fissidentaceae	<i>Grevillea pityophylla</i> 090 Proteaceae
<i>Fissidens taylorii</i> B Fissidentaceae	<i>Grevillea plurijuga</i> 090 Proteaceae
<i>Fissidens vittatus</i> B Fissidentaceae	<i>Grevillea pulchella</i> subsp <i>pulchella</i> ms 090 Proteaceae
<i>Flavoparmelia haysomii</i> L Parmeliaceae	<i>Grevillea rigida</i> subsp <i>rigida</i> 090 Proteaceae
<i>Flavoparmelia rutidota</i> L Parmeliaceae	<i>Grevillea tetrapleura</i> 090 Proteaceae
<i>Fossombronia</i> sp H Hepaticae	<i>Grevillea umbellulata</i> subsp <i>umbellulata</i> 090 Proteaceae
<i>Funaria apophysata</i> B Funariaceae	<i>Grimmia laevigata</i> B Grimmiaceae
<i>Funaria helmsii</i> B Funariaceae	<i>Guichenotia macrantha</i> 223 Sterculiaceae
<i>Funaria products</i> B Funariaceae	<i>Gymnopilus</i> sp F Cortinariaceae
<i>Galerina</i> sp F Cortinariaceae	<i>Gyrostemon subnudus</i> 108 Gyrostemonaceae
<i>Galium aparine</i> 331 Rubiaceae	<i>Haemodorum sparsiflorum</i> 055 Haemodoraceae
<i>Gastrolobium brownii</i> 165 Papilionaceae	<i>Hakea arida</i> 090 Proteaceae
<i>Gastrolobium callistachys</i> 165 Papilionaceae	<i>Hakea clavata</i> 090 Proteaceae
<i>Gastrolobium calycinum</i> 165 Papilionaceae	<i>Hakea erinacea</i> 090 Proteaceae
<i>Gastrolobium involutum</i> 165 Papilionaceae	<i>Hakea myrtoides</i> 090 Proteaceae
<i>Gastrolobium laytonii</i> 165 Papilionaceae	<i>Hakea obliqua</i> 090 Proteaceae
<i>Gastrolobium ovalifolium</i> 165 Papilionaceae	<i>Hakea prostrata</i> 090 Proteaceae
<i>Gastrolobium parviflorum</i> 165 Papilionaceae	<i>Hakea recurva</i> 090 Proteaceae
<i>Gastrolobium spinosum</i> var <i>spinosum</i> 165 Papilionaceae	<i>Hakea suaveolens</i> 090 Proteaceae
<i>Gastrolobium spinosum</i> var <i>trilobum</i> 165 Papilionaceae	<i>Hakea undulata</i> 090 Proteaceae
<i>Genoplesium nigricans</i> 066 Orchidaceae	<i>Haloragodendron glandulosum</i> 276 Haloragaceae
Genus sp 032 Cyperaceae	<i>Haloragodendron racemosum</i> 276 Haloragaceae
Genus sp 185 Euphorbiaceae	<i>Hannalordia bissillii</i> 223 Sterculiaceae
Genus sp Shannon (PG Wilson 1237B) ms 281 Apiaceae	<i>Hannafordia</i> sp 223 Sterculiaceae
<i>Gigaspermum repens</i> B Gigaspermaceae	<i>Heliotropium tenuifolium</i> 310 Boraginaceae
<i>Glischrocaryon aureum</i> var "unsorted" 276 Haloragaceae	<i>Helipterum craspediooides</i> 345 Asteraceae
<i>Glischrocaryon aureum</i> var <i>angustifolium</i> 276 Haloragaceae	<i>Helipterum heteranthum</i> var <i>minor</i> 345 Asteraceae
<i>Glischrocaryon aureum</i> var <i>aureum</i> 276 Haloragaceae	<i>Hemiandra pungens</i> 313 Lamiaceae
<i>Glischrocaryon flavescens</i> 276 Haloragaceae	<i>Hemigenia</i> aff <i>sericea</i> 313 Lamiaceae
<i>Glischrocaryon roei</i> 276 Haloragaceae	<i>Hemigenia incana</i> 313 Lamiaceae
<i>Glischrocaryon</i> sp 276 Haloragaceae	<i>Hemigenia podalyrina</i> 313 Lamiaceae
<i>Glossostigma diandrum</i> 316 Scrophulariaceae	<i>Hemigenia rigidula</i> 313 Lamiaceae
<i>Glossostigma drummondii</i> 316 Scrophulariaceae	<i>Hemigenia sericea</i> 313 Lamiaceae
<i>Glossostigma</i> sp 316 Scrophulariaceae	<i>Hemigenia</i> sp 313 Lamiaceae
<i>Glycine canescens</i> 165 Papilionaceae	<i>Hemigenia</i> sp Albany (GJ Keighery 8712) 313 Lamiaceae
<i>Glycine clandestina</i> 165 Papilionaceae	<i>Hemigenia</i> sp Albany (GJ Keighery 8712) ms 313 Lamiaceae
<i>Gnaphalium indutum</i> 345 Asteraceae	<i>Heterodea muelleri</i> L Heterodeaceae

<i>Heterodermia obscurata</i> L Verrucariaceae	<i>Lampranthus glaucus</i> 110 Aizoaceae
<i>Heteropogon contortus</i> 031 Poaceae	<i>Lasiopetalum compactum</i> 223 Sterculiaceae
<i>Hexagona aparia</i> F Polyporaceae	<i>Lasiopetalum cordifolium</i> 223 Sterculiaceae
<i>Hibbertia acerosa</i> 226 Dilleniaceae	<i>Lawrencella rosea</i> 345 Asteraceae
<i>Hibbertia aff gracilipes</i> (<i>glabrous carpels</i>) 226 Dilleniaceae	<i>Lawrenzia diffusa</i> 221 Malvaceae
<i>Hibbertia aff gracilipes</i> 226 Dilleniaceae	<i>Laxmannia minor</i> 054F Anthericaceae
<i>Hibbertia cunninghamii</i> 226 Dilleniaceae	<i>Laxmannia</i> sp 054F Anthericaceae
<i>Hibbertia glomerosa</i> 226 Dilleniaceae	<i>Laxmannia squarrosa</i> 054F Anthericaceae
<i>Hibbertia gracilipes</i> 226 Dilleniaceae	<i>Lechenaultia formosa</i> 341 Goodeniaceae
<i>Hibbertia granitica</i> 226 Dilleniaceae	<i>Lemooria burkittii</i> 345 Asteraceae
<i>Hibbertia hypericoides</i> 226 Dilleniaceae	<i>Lepidosperma angustatum</i> 032 Cyperaceae
<i>Hibbertia mucronata</i> 226 Dilleniaceae	<i>Lepidosperma brunonianum</i> 032 Cyperaceae
<i>Hibbertia pungens</i> 226 Dilleniaceae	<i>Lepidosperma resinosum</i> 032 Cyperaceae
<i>Hibbertia racemosa</i> 226 Dilleniaceae	<i>Lepidosperma</i> sp 032 Cyperaceae
<i>Hibbertia rhadinopoda</i> 226 Dilleniaceae	<i>Lepidosperma</i> sp A2 Island Flat (Keighery 7000) 032 Cyperaceae
<i>Hibbertia rupicola</i> 226 Dilleniaceae	<i>Lepidosperma</i> sp K 032 Cyperaceae
<i>Hibbertia spicata</i> 226 Dilleniaceae	<i>Lepidosperma tuberculatum</i> 032 Cyperaceae
<i>Hibiscus leptocladius</i> 221 Malvaceae	<i>Lepiota</i> sp F Agaricaceae
<i>Homalosciadium homalocarpum</i> 281 Apiaceae	<i>Leporella fimbriata</i> 066 Orchidaceae
<i>Hovea acanthoclada</i> 165 Papilionaceae	<i>Leprocaulon microscopicum</i> L <i>Fungi Imperfecti</i>
<i>Hovea pungens</i> 165 Papilionaceae	<i>Leproloma membranaceum</i> L ?Pannariaceae
<i>Hyalochlamys globifera</i> 345 Asteraceae	<i>Leptocarpus</i> aff <i>roycei</i> "white" 039 Restionaceae
<i>Hyalosperma glutinosum</i> subsp <i>glutinosum</i> 345 Asteraceae	<i>Leptocarpus kraussii</i> ms 039 Restionaceae
<i>Hyalosperma pusillum</i> 345 Asteraceae	<i>Leptocarpus</i> <i>roycei</i> ms 039 Restionaceae
<i>Hybanthus floribundus</i> subsp <i>floribundus</i> 243 Violaceae	<i>Leptoceras menziesii</i> 066 Orchidaceae
<i>Hybanthus monopetalus</i> 243 Violaceae	<i>Leptodontium paradoxum</i> B Pottiaceae
<i>Hydrocotyle alata</i> 281 Apiaceae	<i>Leptomeria cunninghamii</i> 092 Santalaceae
<i>Hydrocotyle callicarpa</i> 281 Apiaceae	<i>Leptomeria squarrulosa</i> 092 Santalaceae
<i>Hydrocotyle diantha</i> 281 Apiaceae	<i>Leptosema aphyllum</i> ms 165 Papilionaceae
<i>Hydrocotyle puberula</i> ms 281 Apiaceae	<i>Leptospermum</i> aff <i>roei</i> 273 Myrtaceae
<i>Hydrocotyle scutellifera</i> 281 Apiaceae	<i>Leptospermum erubescens</i> 273 Myrtaceae
<i>Hypericum japonicum</i> 233 Clusiaceae	<i>Leptospermum incanum</i> 273 Myrtaceae
<i>Hypocalymma angustifolium</i> 273 Myrtaceae	<i>Leptospermum macgillivrayi</i> 273 Myrtaceae
<i>Hypocalymma uncinatum</i> ms 273 Myrtaceae	<i>Leptospermum roei</i> 273 Myrtaceae
<i>Hypochoeris glabra</i> 345 Asteraceae	<i>Leptospermum sericeum</i> 273 Myrtaceae
<i>Hypogymnia subphysodes</i> L Hypogymniaceae	<i>Leptospermum</i> sp 273 Myrtaceae
<i>Hypoxis glabella</i> var <i>glabella</i> 056A Hypoxidaceae	<i>Lethocolea squamata</i> H Hepaticae
<i>Hypoxis occidentalis</i> var <i>quadriloba</i> 056A Hypoxidaceae	<i>Leucopogon</i> aff <i>reflexus</i> 288 Epacridaceae
<i>Indigofera australis</i> 165 Papilionaceae	<i>Leucopogon australis</i> 288 Epacridaceae
<i>Indigofera monophylla</i> 165 Papilionaceae	<i>Leucopogon bracteolaris</i> 288 Epacridaceae
<i>Ipomoea nil</i> 307 Convolvulaceae	<i>Leucopogon glabellus</i> 288 Epacridaceae
<i>Ischyrodon lepturus</i> B Brachytheciaceae	<i>Leucopogon pendulus</i> 288 Epacridaceae
<i>Isoetes australis</i> 004 Isoetaceae	<i>Leucopogon reflexus</i> 288 Epacridaceae
<i>Isoetes caroli</i> 004 Isoetaceae	<i>Leucopogon revolutus</i> 288 Epacridaceae
<i>Isoetes inflata</i> 004 Isoetaceae	<i>Leucopogon rotundifolius</i> 288 Epacridaceae
<i>Isoetes</i> sp 004 Isoetaceae	<i>Leucopogon</i> sp 288 Epacridaceae
<i>Isolepis congrua</i> 032 Cyperaceae	<i>Leucopogon sprengeliooides</i> 288 Epacridaceae
<i>Isolepis marginata</i> 032 Cyperaceae	<i>Leucopogon strictus</i> 288 Epacridaceae
<i>Isopogon attenuatus</i> 090 Proteaceae	<i>Leucopogon unilateralis</i> 288 Epacridaceae
<i>Isopogon divergens</i> 090 Proteaceae	<i>Levenhookia dubia</i> 343 Stylidiaceae
<i>Isopogon formosus</i> 090 Proteaceae	<i>Levenhookia leptantha</i> 343 Stylidiaceae
<i>Isotoma hypocrateiformis</i> 340 Lobeliaceae	<i>Levenhookia pauciflora</i> 343 Stylidiaceae
<i>Isotoma petraea</i> 340 Lobeliaceae	<i>Levenhookia pusilla</i> 343 Stylidiaceae
<i>Isotoma scapigera</i> 340 Lobeliaceae	<i>Levenhookia stipitata</i> 343 Stylidiaceae
<i>Isotropis atropurpurea</i> 165 Papilionaceae	<i>Lobelia alata</i> 340 Lobeliaceae
<i>Isotropis cuneifolia</i> 165 Papilionaceae	<i>Lobelia gibbosa</i> 340 Lobeliaceae
<i>Isotropis</i> sp 165 Papilionaceae	<i>Lobelia heterophylla</i> 340 Lobeliaceae
<i>Jacksonia alata</i> 165 Papilionaceae	<i>Lobelia</i> sp 340 Lobeliaceae
<i>Jacksonia furcellata</i> 165 Papilionaceae	<i>Logania serpyllifolia</i> subsp <i>serpyllifolia</i> 302 Loganiaceae
<i>Jacksonia spinosa</i> 165 Papilionaceae	<i>Logania</i> sp 302 Loganiaceae
<i>Jamesoniella colorata</i> H Hepaticae	<i>Logania stenophylla</i> 302 Loganiaceae
<i>Juncus bufonius</i> 052 Juncaceae	<i>Lolium perenne</i> 031 Poaceae
<i>Juncus capitatus</i> 052 Juncaceae	<i>Lomandra integrifolia</i> 054C Dasypogonaceae
<i>Juncus pauciflorus</i> 052 Juncaceae	<i>Lomandra rigidula</i> 054C Dasypogonaceae
<i>Kennedia beckxiana</i> 165 Papilionaceae	<i>Lotus angustissimus</i> 165 Papilionaceae
<i>Kennedia carinata</i> 165 Papilionaceae	<i>Lycoperdon</i> sp F Lycoperdaceae
<i>Kennedia glabrata</i> 165 Papilionaceae	<i>Lyperanthus serratus</i> 066 Orchidaceae
<i>Kennedia macrophylla</i> 165 Papilionaceae	<i>Macrozamia riedlei</i> 016A Zamiaceae
<i>Kennedia prostrata</i> 165 Papilionaceae	<i>Maireana glomerifolia</i> 105 Chenopodiaceae
<i>Kennedia stirlingii</i> 165 Papilionaceae	<i>Malleostemon tuberculatus</i> 273 Myrtaceae
<i>Keraudrenia integrifolia</i> 223 Sterculiaceae	<i>Marsdenia granitica</i> 305 Asclepiadaceae
<i>Kunzea aff pulchella</i> 273 Myrtaceae	<i>Marsilea drummondii</i> 013 Marsileaceae
<i>Kunzea affinis</i> 273 Myrtaceae	<i>Melaleuca adnata</i> 273 Myrtaceae
<i>Kunzea baxteri</i> 273 Myrtaceae	<i>Melaleuca ctenoides</i> 273 Myrtaceae
<i>Kunzea micromera</i> 273 Myrtaceae	<i>Melaleuca cuticularis</i> 273 Myrtaceae
<i>Kunzea pulchella</i> 273 Myrtaceae	<i>Melaleuca diosmifolia</i> 273 Myrtaceae
<i>Labichea lanceolata</i> subsp <i>brevifolia</i> 164 Caesalpiniaceae	<i>Melaleuca eleuterostachya</i> 273 Myrtaceae
<i>Labichea teretifolia</i> subsp <i>grandistipulata</i> 164 Caesalpiniaceae	<i>Melaleuca elliptica</i> 273 Myrtaceae
<i>Laccaria laccata</i> F Tricholomataceae	<i>Melaleuca eximia</i> ms 273 Myrtaceae
<i>Lambertia inermis</i> var <i>inermis</i> 090 Proteaceae	<i>Melaleuca fulgens</i> subsp <i>fulgens</i> 273 Myrtaceae

<i>Melaleuca fulgens</i> subsp <i>steedmanii</i> 273 Myrtaceae	<i>Patersonia occidentalis</i> 060 Iridaceae
<i>Melaleuca glaberrima</i> 273 Myrtaceae	<i>Pelargonium australe</i> 167 Geraniaceae
<i>Melaleuca globifera</i> 273 Myrtaceae	<i>Pelargonium</i> cf. <i>littorale</i> 167 Geraniaceae
<i>Melaleuca hamulosa</i> 273 Myrtaceae	<i>Pelargonium drummondii</i> 167 Geraniaceae
<i>Melaleuca holosericea</i> 273 Myrtaceae	<i>Pelargonium hylasae</i> 167 Geraniaceae
<i>Melaleuca lasiandra</i> 273 Myrtaceae	<i>Pelargonium</i> sp 167 Geraniaceae
<i>Melaleuca laxiflora</i> 273 Myrtaceae	<i>Pentaschistis airoides</i> 031 Poaceae
<i>Melaleuca leiocarpa</i> 273 Myrtaceae	<i>Pericalymma ellipticum</i> var <i>floridum</i> ms 273 Myrtaceae
<i>Melaleuca macronychia</i> subsp <i>macronychia</i> 273 Myrtaceae	<i>Persoonia amaliae</i> 090 Proteaceae
<i>Melaleuca pentagona</i> var <i>subulifolia</i> 273 Myrtaceae	<i>Persoonia falcata</i> 090 Proteaceae
<i>Melaleuca pulchella</i> 273 Myrtaceae	<i>Persoonia pentasticha</i> ms 090 Proteaceae
<i>Melaleuca pungens</i> var "unsorted" 273 Myrtaceae	<i>Persoonia quinquenervis</i> 090 Proteaceae
<i>Melaleuca radula</i> 273 Myrtaceae	<i>Petalostylis cassioides</i> 164 Caesalpiniaceae
<i>Melaleuca scabra</i> 273 Myrtaceae	<i>Petrophile divaricata</i> 090 Proteaceae
<i>Melaleuca</i> sp 273 Myrtaceae	<i>Petrophile squamata</i> 090 Proteaceae
<i>Melaleuca uncinata</i> 273 Myrtaceae	<i>Petrophile striata</i> 090 Proteaceae
<i>Melaleuca urceolaris</i> 273 Myrtaceae	<i>Petrorrhagia velutina</i> 113 Caryophyllaceae
<i>Melaleuca viminea</i> 273 Myrtaceae	<i>Phedalium elegans</i> ms 175 Rutaceae
<i>Melilotus indicus</i> 165 Papilionaceae	<i>Phedalium filfolium</i> 175 Rutaceae
<i>Menegazzia foraminulosa</i> L Hypogynniamiaeae	<i>Phedalium rhytidophyllum</i> 175 Rutaceae
<i>Menegazzia subpertusa</i> L Hypogynniamiaeae	<i>Philothea langei</i> 175 Rutaceae
<i>Mentha spicata</i> 313 Lamiaceae	<i>Philydrella pygmaea</i> subsp "unsorted" 050 Philydraceae
<i>Mesomelaena tetragona</i> 032 Cyperaceae	<i>Philydrella pygmaea</i> subsp <i>pygmaea</i> 050 Philydraceae
<i>Micromyrtus serrulata</i> 273 Myrtaceae	<i>Phyllangium divergens</i> 302 Loganiaceae
<i>Micromyrtus sulphurea</i> 273 Myrtaceae	<i>Phyllangium paradoxum</i> ms 302 Loganiaceae
<i>Microtis aff parviflora</i> 066 Orchidaceae	<i>Phyllanthus calycinus</i> 185 Euphorbiaceae
<i>Microtis atrata</i> 066 Orchidaceae	<i>Phyllanthus</i> sp 185 Euphorbiaceae
<i>Microtis brownii</i> 066 Orchidaceae	<i>Pimelea brachyphylla</i> 263 Thymelaeaceae
<i>Microtis media</i> subsp "inland race" 066 Orchidaceae	<i>Pimelea ciliata</i> 263 Thymelaeaceae
<i>Microtis media</i> subsp <i>media</i> 066 Orchidaceae	<i>Pimelea ferruginea</i> 263 Thymelaeaceae
<i>Microtis rara</i> 066 Orchidaceae	<i>Pimelea forrestiana</i> 263 Thymelaeaceae
<i>Millotia depauperata</i> 345 Asteraceae	<i>Pimelea graniticola</i> 263 Thymelaeaceae
<i>Millotia myosotidifolia</i> 345 Asteraceae	<i>Pimelea imbricata</i> var <i>imbricata</i> 263 Thymelaeaceae
<i>Millotia tenuifolia</i> var <i>tenuifolia</i> 345 Asteraceae	<i>Pimelea imbricata</i> var <i>pilgera</i> 263 Thymelaeaceae
<i>Mirbelia longifolia</i> 165 Papilionaceae	<i>Pimelea microcephala</i> subsp <i>microcephala</i> 263 Thymelaeaceae
<i>Mirbelia</i> sp 165 Papilionaceae	<i>Pimelea preissii</i> 263 Thymelaeaceae
<i>Mitrasacme nudicaulis</i> 302 Loganiaceae	<i>Pimelea</i> sp 263 Thymelaeaceae
<i>Mitrasacme</i> sp 302 Loganiaceae	<i>Pimelea spiculigera</i> var <i>thesioides</i> 263 Thymelaeaceae
<i>Monadenia bracteata</i> 066 Orchidaceae	<i>Pimelea suaveolens</i> subsp <i>suaveolens</i> 263 Thymelaeaceae
<i>Monotaxis grandiflora</i> 185 Euphorbiaceae	<i>Pimelea sulphurea</i> 263 Thymelaeaceae
<i>Muehlenbeckia adpressa</i> 103 Polygonaceae	<i>Pittosporum phylliraeoides</i> 152 Pittosporaceae
<i>Muelleranthus trifoliolatus</i> 165 Papilionaceae	<i>Pityrodia dilatata</i> 311A Chloanthaceae
<i>Mukia maderaspatana</i> 337 Cucurbitaceae	<i>Pityrodia teckiana</i> 311A Chloanthaceae
<i>Myriocephalus guerinae</i> 345 Asteraceae	<i>Pityrodia terminalis</i> 311A Chloanthaceae
<i>Myriocephalus nudus</i> 345 Asteraceae	<i>Plantago debilis</i> 329 Plantaginaceae
<i>Myriocephalus occidentalis</i> 345 Asteraceae	<i>Platysace compressa/filiformis</i> 281 Apiaceae
<i>Myriocephalus pygmaeus</i> 345 Asteraceae	<i>Platysace compressa</i> 281 Apiaceae
<i>Myriophyllum balladoniense</i> 276 Haloragaceae	<i>Pleuridium nervosum</i> B Ditrichaceae
<i>Myriophyllum lapidicola</i> 276 Haloragaceae	<i>Pleurosorus rutifolius</i> 011E Aspleniaceae
<i>Myriophyllum petraeum</i> 276 Haloragaceae	<i>Pleurosorus subglandulosus</i> 011E Aspleniaceae
<i>Myriophyllum</i> sp 276 Haloragaceae	<i>Pleurotus nidiformis</i> F Polyporaceae
<i>Nelsonia campestris</i> 325 Acanthaceae	<i>Poa annua</i> 031 Poaceae
<i>Nemcia acuta</i> 165 Papilionaceae	<i>Poa poiformis</i> 031 Poaceae
<i>Neofuscelia pulla</i> L Parmeliaceae	<i>Podolepis canescens</i> 345 Asteraceae
<i>Neurachne alopecuroides</i> 031 Poaceae	<i>Podolepis capillaris</i> 345 Asteraceae
<i>Nicotiana cavicola</i> 315 Solanaceae	<i>Podolepis lessonii</i> 345 Asteraceae
<i>Nicotiana occidentalis</i> subsp <i>obliqua</i> 315 Solanaceae	<i>Podolepis rugata</i> 345 Asteraceae
<i>Nicotiana rotundifolia</i> 315 Solanaceae	<i>Podotheca angustifolia</i> 345 Asteraceae
<i>Olearia aff paucidentata</i> 345 Asteraceae	<i>Podotheca wilsonii</i> 345 Asteraceae
<i>Olearia algida</i> 345 Asteraceae	<i>Pogonolepis stricta</i> 345 Asteraceae
<i>Olearia muelleri</i> 345 Asteraceae	<i>Polycarpa breviflora</i> var <i>gracilis</i> 113 Caryophyllaceae
<i>Olearia paucidentata</i> 345 Asteraceae	<i>Polycarpa corymbosa</i> var "unsorted" 113 Caryophyllaceae
<i>Olearia propinqua</i> 345 Asteraceae	<i>Polygala orbicularis</i> 183 Polygalaceae
<i>Olearia stuartii</i> 345 Asteraceae	<i>Porana commixta</i> 307 Convolvulaceae
<i>Omphalappula concava</i> 310 Boraginaceae	<i>Porana sericea</i> 307 Convolvulaceae
<i>Opercularia vaginalis</i> 331 Rubiaceae	<i>Pottia drummondii</i> B Pottiaceae
<i>Ophioglossum lusitanicum</i> 005 Ophioglossaceae	<i>Prasophyllum aff parvifolium</i> 066 Orchidaceae
<i>Ophioglossum</i> sp 005 Ophioglossaceae	<i>Prasophyllum brownii</i> 066 Orchidaceae
<i>Osteospermum clandestinum</i> 345 Asteraceae	<i>Prasophyllum cucullatum</i> 066 Orchidaceae
<i>Oudemansiella</i> sp F Tricholomataceae	<i>Prasophyllum elatum</i> 066 Orchidaceae
<i>Ozothamnus alpinus</i> 345 Asteraceae	<i>Prasophyllum fimbria</i> 066 Orchidaceae
<i>Ozothamnus ramosus</i> 345 Asteraceae	<i>Prasophyllum gibbosum</i> 066 Orchidaceae
<i>Pandorea pandorana</i> 317 Bignoniaceae	<i>Prasophyllum gibbosum</i> subsp <i>gibbosum</i> 066 Orchidaceae
<i>Paracaleana nigrita</i> 066 Orchidaceae	<i>Prasophyllum gracile</i> 066 Orchidaceae
<i>Paracaleana triens</i> ms 066 Orchidaceae	<i>Prasophyllum parvifolium</i> 066 Orchidaceae
<i>Paraserianthes lophantha</i> subsp <i>lophantha</i> 163 Mimosaceae	<i>Prasophyllum ringens</i> 066 Orchidaceae
<i>Parentucellia latifolia</i> 316 Scrophulariaceae	<i>Prasophyllum</i> sp 066 Orchidaceae
<i>Parentucellia viscosa</i> 316 Scrophulariaceae	<i>Prostanthera baxteri</i> 313 Lamiaceae
<i>Parmelia</i> sp L Parmeliaceae	<i>Prostanthera campbellii</i> 313 Lamiaceae
<i>Paspalidium clementii</i> 031 Poaceae	<i>Prostanthera carrickiana</i> 313 Lamiaceae

<i>Prostanthera florifera</i> 313 Lamiaceae	<i>Sclerolaena costata</i> 105 Chenopodiaceae
<i>Prostanthera gryloana</i> 313 Lamiaceae	<i>Sclerolaena fusiformis</i> 105 Chenopodiaceae
<i>Prostanthera magnifica</i> 313 Lamiaceae	<i>Scyphocoris major</i> 345 Asteraceae
<i>Prostanthera serpyllifolia</i> subsp <i>microphylla</i> 313 Lamiaceae	<i>Selaginella gracillima</i> 003 Selaginellaceae
<i>Prostanthera</i> sp 1 313 Lamiaceae	<i>Sematophyllum homomallum</i> B Sematophyllaceae
<i>Prostanthera striatiflora</i> 313 Lamiaceae	<i>Senecio hispidulus</i> var <i>hispidulus</i> 345 Asteraceae
<i>Prostanthera verticillaris</i> 313 Lamiaceae	<i>Senecio leucoglossus</i> 345 Asteraceae
<i>Pterochaeta paniculata</i> 345 Asteraceae	<i>Senna artemisioides</i> subsp <i>x artemisioides</i> 164 Caesalpiniaceae
<i>Pterostylis aff nana</i> 066 Orchidaceae	<i>Senna artemisioides</i> subsp <i>x sturtii</i> 164 Caesalpiniaceae
<i>Pterostylis aff rufa</i> 066 Orchidaceae	<i>Senna glutinosa</i> subsp <i>charlesiana</i> 164 Caesalpiniaceae
<i>Pterostylis allantoidea</i> 066 Orchidaceae	<i>Sida calyxhymenia</i> 221 Malvaceae
<i>Pterostylis aspera</i> 066 Orchidaceae	<i>Sida hookeriana</i> 221 Malvaceae
<i>Pterostylis barbata</i> 066 Orchidaceae	<i>Siloxerus filifolius</i> 345 Asteraceae
<i>Pterostylis elegans</i> 066 Orchidaceae	<i>Siphula coriacea</i> L Siphonaceae
<i>Pterostylis hamiltonii</i> 066 Orchidaceae	<i>Sisymbrium irio</i> 138 Brassicaceae
<i>Pterostylis mutica</i> 066 Orchidaceae	<i>Solanum cleistogamum</i> 315 Solanaceae
<i>Pterostylis recurva</i> 066 Orchidaceae	<i>Solanum ferociissimum</i> 315 Solanaceae
<i>Pterostylis roensis</i> 066 Orchidaceae	<i>Solanum lasiophyllum</i> 315 Solanaceae
<i>Pterostylis sanguinea</i> 066 Orchidaceae	<i>Solanum lucani</i> 315 Solanaceae
<i>Pterostylis sargentii</i> 066 Orchidaceae	<i>Solanum nummularium</i> 315 Solanaceae
<i>Pterostylis scabra</i> 066 Orchidaceae	<i>Solanum orbiculatum</i> subsp <i>orbiculatum</i> 315 Solanaceae
<i>Pterostylis vittata</i> 066 Orchidaceae	<i>Solanum phlomoides</i> 315 Solanaceae
<i>Ptilotus corymbosus</i> var <i>corymbosus</i> 106 Amaranthaceae	<i>Solanum symonii</i> 315 Solanaceae
<i>Ptilotus divaricatus</i> var "unsorted" 106 Amaranthaceae	<i>Sollya heterophylla</i> 152 Pittosporaceae
<i>Ptilotus gaudichaudii</i> var "unsorted" 106 Amaranthaceae	<i>Sonchus asper</i> sens lat 345 Asteraceae
<i>Ptilotus gaudichaudii</i> var <i>parviflorus</i> 106 Amaranthaceae	<i>Spartochloa scirpoidea</i> 031 Poaceae
<i>Ptilotus humilis</i> var <i>humilis</i> 106 Amaranthaceae	<i>Spergularia rubra</i> 113 Caryophyllaceae
<i>Ptilotus incanus</i> var <i>incanus</i> 106 Amaranthaceae	<i>Sphenotoma capitatum</i> 288 Epacridaceae
<i>Ptilotus obovatus</i> var "unsorted" 106 Amaranthaceae	<i>Spiculaea ciliata</i> 066 Orchidaceae
<i>Ptilotus obovatus</i> var <i>obovatus</i> 106 Amaranthaceae	<i>Stackhousea monogyna</i> 202 Stackhousiaceae
<i>Ptilotus polystachys</i> forma <i>rubriflorus</i> 106 Amaranthaceae	<i>Stackhousea muricata</i> 202 Stackhousiaceae
<i>Ptilotus spathulatus</i> forma <i>spathulatus</i> 106 Amaranthaceae	<i>Stenantherum notiale</i> subsp <i>notiale</i> 215 Rhamnaceae
<i>Pultenaea elachista</i> 165 Papilionaceae	<i>Stipa compressa</i> 031 Poaceae
<i>Pultenaea ericifolia</i> 165 Papilionaceae	<i>Stipa hemipogon</i> 031 Poaceae
<i>Pultenaea obcordata</i> 165 Papilionaceae	<i>Stipa variabilis</i> 031 Poaceae
<i>Punctelia subrudecta</i> L Parmeliaceae	<i>Stropharia</i> sp <i>F Strophariaceae</i>
<i>Pyrorchis nigricans</i> 066 Orchidaceae	<i>Styliodium amoenum</i> var "unsorted" 343 Stylidiaceae
<i>Quinetia urvillei</i> 345 Asteraceae	<i>Styliodium arenicola</i> 343 Stylidiaceae
<i>Quinqueremulus linearis</i> 345 Asteraceae	<i>Styliodium assimile</i> 343 Stylidiaceae
<i>Ranunculus colonorum</i> 119 Ranunculaceae	<i>Styliodium breviscapum</i> var <i>breviscapum</i> 343 Stylidiaceae
<i>Ranunculus</i> sp 119 Ranunculaceae	<i>Styliodium brunonianum</i> subsp "unsorted" 343 Stylidiaceae
<i>Restio crispatus</i> 039 Restionaceae	<i>Styliodium brunonianum</i> subsp <i>brunonianum</i> 343 Stylidiaceae
<i>Rhadinothamnus euphemiae</i> 175 Rutaceae	<i>Styliodium bulbiferum</i> var "unsorted" 343 Stylidiaceae
<i>Rhagodia eremaea</i> 105 Chenopodiaceae	<i>Styliodium caespitosum</i> 343 Stylidiaceae
<i>Rhagodia preissii</i> subsp <i>preissii</i> 105 Chenopodiaceae	<i>Styliodium calcaratum</i> 343 Stylidiaceae
<i>Rhodanthe battii</i> 345 Asteraceae	<i>Styliodium crassifolium</i> 343 Stylidiaceae
<i>Rhodanthe chlorocephala</i> subsp <i>splendida</i> 345 Asteraceae	<i>Styliodium despactum</i> 343 Stylidiaceae
<i>Rhodanthe citrina</i> 345 Asteraceae	<i>Styliodium dichotomum</i> 343 Stylidiaceae
<i>Rhodanthe frenchii</i> 345 Asteraceae	<i>Styliodium dielsianum</i> 343 Stylidiaceae
<i>Rhodanthe manglesii</i> 345 Asteraceae	<i>Styliodium divaricatum</i> 343 Stylidiaceae
<i>Rhodanthe rubella</i> 345 Asteraceae	<i>Styliodium diversifolium</i> 343 Stylidiaceae
<i>Rhodanthe spicata</i> 345 Asteraceae	<i>Styliodium ecorne</i> 343 Stylidiaceae
<i>Rhodanthe tietkensii</i> 345 Asteraceae	<i>Styliodium emarginatum</i> subsp <i>emarginatum</i> 343 Stylidiaceae
<i>Riccia crinita</i> H Ricciaceae	<i>Styliodium inundatum</i> 343 Stylidiaceae
<i>Riccia crystallina</i> H Ricciaceae	<i>Styliodium longibracteatum</i> 343 Stylidiaceae
<i>Riccia lamellosa</i> H Ricciaceae	<i>Styliodium merrallii</i> 343 Stylidiaceae
<i>Riccia limbata</i> H Ricciaceae	<i>Styliodium neglectum</i> 343 Stylidiaceae
<i>Ricinocarpus glaucus</i> 185 Euphorbiaceae	<i>Styliodium nungarinense</i> 343 Stylidiaceae
<i>Ricinocarpus rosmarinifolius</i> 185 Euphorbiaceae	<i>Styliodium perpusillum</i> 343 Stylidiaceae
<i>Rimelia reticulata</i> L Parmeliaceae	<i>Styliodium pulchellum</i> 343 Stylidiaceae
<i>Rinzia</i> sp 273 Myrtaceae	<i>Styliodium pygmaeum</i> 343 Stylidiaceae
<i>Rulingia crauophylla</i> 223 Sterculiaceae	<i>Styliodium sp</i> 343 Stylidiaceae
<i>Rulingia cygnorum</i> 223 Sterculiaceae	<i>Styliodium spathulatum</i> subsp "unsorted" 343 Stylidiaceae
<i>Rulingia kempeana</i> 223 Sterculiaceae	<i>Styliodium spathulatum</i> subsp <i>acuminatum</i> 343 Stylidiaceae
<i>Rulingia luteiflora</i> 223 Sterculiaceae	<i>Stypandra glauca</i> 054E Phormiaceae
<i>Russula</i> "small white" (K Syme 633/93) F Russulaceae	<i>Swainsona gracilis</i> 165 Papilionaceae
<i>Rutidosis multiflora</i> 345 Asteraceae	<i>Swainsona</i> sp 165 Papilionaceae
<i>Samolus repens</i> subsp "unsorted" 293 Primulaceae	<i>Synaphea</i> sp 090 Proteaceae
<i>Santalum spicatum</i> 092 Santalaceae	<i>Templetonia sulcata</i> 165 Papilionaceae
<i>Scaevela glandulifera</i> 341 Goodeniaceae	<i>Tephrosia</i> sp 165 Papilionaceae
<i>Scaevela spinescens</i> 341 Goodeniaceae	<i>Tephrosia</i> sp <i>B Kimberley Flora</i> (CA Gardner 7) 165 Papilionaceae
<i>Scaevela thesioides</i> subsp <i>filifolia</i> 341 Goodeniaceae	<i>Tetragonia cristata</i> 110 Aizoaceae
<i>Schizaea</i> sp 006 Schizaeaceae	<i>Tetrapterum cylindricum</i> B Pottiaceae
<i>Schoenia ayersii</i> 345 Asteraceae	<i>Tetrapeltis capillaris</i> 032 Cyperaceae
<i>Schoenia cassiniiana</i> 345 Asteraceae	<i>Tetratheca aff harperi</i> 182 Tremandraceae
<i>Schoenus</i> aff <i>odontocarpus</i> 032 Cyperaceae	<i>Tetratheca deltoidea</i> 182 Tremandraceae
<i>Schoenus multiglumis</i> 032 Cyperaceae	<i>Tetratheca hirsuta</i> 182 Tremandraceae
<i>Schoenus odontocarpus</i> (Small typical variant) 032 Cyperaceae	<i>Tetratheca hispidissima</i> 182 Tremandraceae
<i>Schoenus sculptus</i> 032 Cyperaceae	<i>Tetratheca nuda</i> 182 Tremandraceae
<i>Schoenus</i> sp 3 (aff <i>odontocarpus</i>) 032 Cyperaceae	<i>Tetratheca paynterae</i> 182 Tremandraceae

<i>Thelymitra</i> SP 066 Orchidaceae	<i>Triquetrella papillata</i> B Pottiaceae
<i>Thelymitra</i> aff <i>hoelmsii</i> 066 Orchidaceae	<i>Trymalium ledifolium</i> var <i>rosmarinifolium</i> 215 Rhamnaceae
<i>Thelymitra</i> aff <i>longifolia</i> 066 Orchidaceae	<i>Trymalium spatulatum</i> 215 Rhamnaceae
<i>Thelymitra</i> aff <i>nuda</i> 066 Orchidaceae	<i>Tylophora</i> sp 305 Asclepiadaceae
<i>Thelymitra</i> aff <i>pauciflora</i> 066 Orchidaceae	<i>Ursinia anthemoides</i> 345 Asteraceae
<i>Thelymitra</i> <i>antennifera</i> 066 Orchidaceae	<i>Usnea confusa</i> L Parmeliaceae
<i>Thelymitra</i> <i>benthamiana</i> 066 Orchidaceae	<i>Usnea inermis</i> L Usneaceae
<i>Thelymitra</i> <i>crinita</i> 066 Orchidaceae	<i>Utricularia menziesii</i> 323 Lentibulariaceae
<i>Thelymitra</i> <i>cincinnata</i> 066 Orchidaceae	<i>Utricularia multifida</i> 323 Lentibulariaceae
<i>Thelymitra</i> <i>dedmaniarum</i> 066 Orchidaceae	<i>Utricularia violacea</i> 323 Lentibulariaceae
<i>Thelymitra</i> <i>flexuosa</i> 066 Orchidaceae	<i>Velleia cycnopotamica</i> 341 Goodeniaceae
<i>Thelymitra</i> <i>macmillanii</i> 066 Orchidaceae	<i>Velleia trinervis</i> 341 Goodeniaceae
<i>Thelymitra</i> <i>macrophylla</i> 066 Orchidaceae	<i>Verticordia acerosa</i> var <i>acerosa</i> 273 Myrtaceae
<i>Thelymitra</i> <i>nuda</i> 066 Orchidaceae	<i>Verticordia acerosa</i> var <i>preissii</i> 273 Myrtaceae
<i>Thelymitra</i> sp 066 Orchidaceae	<i>Verticordia brownii</i> 273 Myrtaceae
<i>Thelymitra</i> sp 066 Orchidaceae	<i>Verticordia chrysanthae</i> 273 Myrtaceae
<i>Thelymitra</i> <i>spiralis</i> 066 Orchidaceae	<i>Verticordia chrysanthella</i> 273 Myrtaceae
<i>Themeda</i> <i>triandra</i> 031 Poaceae	<i>Verticordia endlicheriana</i> var <i>angustifolia</i> 273 Myrtaceae
<i>Thomasia</i> <i>foliosa</i> 223 Sterculiaceae	<i>Verticordia habrantha</i> 273 Myrtaceae
<i>Thomasia</i> <i>glutinosa</i> var <i>latifolia</i> 223 Sterculiaceae	<i>Verticordia helmsii</i> 273 Myrtaceae
<i>Thomasia</i> <i>grandiflora</i> 223 Sterculiaceae	<i>Verticordia huegelii</i> var <i>decumbens</i> 273 Myrtaceae
<i>Thomasia</i> <i>multiflora</i> 223 Sterculiaceae	<i>Verticordia huegelii</i> var <i>huegelii</i> 273 Myrtaceae
<i>Thomasia</i> <i>pauciflora</i> 223 Sterculiaceae	<i>Verticordia huegelii</i> var <i>tridens</i> 273 Myrtaceae
<i>Thomasia</i> <i>petalocalyx</i> 223 Sterculiaceae	<i>Verticordia minutiflora</i> 273 Myrtaceae
<i>Thomasia</i> sp 223 Sterculiaceae	<i>Verticordia penicillaris</i> 273 Myrtaceae
<i>Thryptomene</i> <i>australis</i> 273 Myrtaceae	<i>Verticordia pennigera</i> 273 Myrtaceae
<i>Thryptomene</i> <i>muconulata</i> 273 Myrtaceae	<i>Verticordia plumosa</i> 273 Myrtaceae
<i>Thryptomene</i> <i>saxicola</i> 273 Myrtaceae	<i>Verticordia plumosa</i> var <i>grandiflora</i> 273 Myrtaceae
<i>Thryptomene</i> sp 273 Myrtaceae	<i>Verticordia plumosa</i> var <i>plumosa</i> 273 Myrtaceae
<i>Thuidium</i> <i>sparsum</i> var <i>hastatum</i> B Thuidiaceae	<i>Verticordia staminosa</i> subsp <i>cylindracea</i> var <i>cylindracea</i> 273 Myrtaceae
<i>Thysanothecium</i> <i>hookeri</i> L Cladoniaceae	<i>Verticordia staminosa</i> subsp <i>cylindracea</i> var <i>erecta</i> 273 Myrtaceae
<i>Thysanotus</i> <i>dichotomus</i> 054F Anthericaceae	<i>Verticordia staminosa</i> subsp <i>staminosa</i> 273 Myrtaceae
<i>Thysanotus</i> <i>manglesianus</i> 054F Anthericaceae	<i>Viminaria juncea</i> 165 Papilionaceae
<i>Thysanotus</i> <i>sparteus</i> 054F Anthericaceae	<i>Vulpia myuros</i> 031 Poaceae
<i>Thysanotus</i> <i>tenellus</i> 054F Anthericaceae	<i>Wahlenbergia preissii</i> 339 Campanulaceae
<i>Thysanotus</i> <i>triandrus</i> 054F Anthericaceae	<i>Waitzia acuminata</i> var <i>acuminata</i> 345 Asteraceae
<i>Tinospora</i> <i>smilacina</i> 122 Menispermaceae	<i>Waitzia nitida</i> 345 Asteraceae
<i>Tolpis</i> <i>barbata</i> 345 Asteraceae	<i>Waitzia podolepis</i> 345 Asteraceae
<i>Tortula</i> <i>antarctica</i> B Pottiaceae	<i>Waitzia suaveolens</i> var <i>flava</i> 345 Asteraceae
<i>Tortula</i> <i>principis</i> B Pottiaceae	<i>Weissia brachycarpa</i> B Pottiaceae
<i>Trachymene</i> <i>cyanopetala</i> 281 Apiaceae	<i>Weissia controversa</i> B Pottiaceae
<i>Trachymene</i> <i>moorei</i> subsp <i>Tutanning</i> (AS George 12867) ms 281 Apiaceae	<i>Weissia rutilans</i> B Pottiaceae
<i>Trachymene</i> <i>ornata</i> 281 Apiaceae	<i>Westringia dampieri</i> 313 Lamiaceae
<i>Trachymene</i> <i>pilosa</i> 281 Apiaceae	<i>Wurmbea cernua</i> 054J Colchicaceae
<i>Trachymene</i> sp <i>Pilbara</i> (RA Saffrey 1117) ms 281 Apiaceae	<i>Wurmbea densiflora</i> 054J Colchicaceae
<i>Tremandra</i> <i>diffusa</i> 182 Tremandraceae	<i>Wurmbea dioica</i> 054J Colchicaceae
<i>Tremandra</i> <i>stelligera</i> 182 Tremandraceae	<i>Wurmbea dioica</i> subsp <i>alba</i> 054J Colchicaceae
<i>Tribonanthes</i> <i>australis</i> 055 Haemodoraceae	<i>Wurmbea murchisoniana</i> 054J Colchicaceae
<i>Tribonanthes</i> <i>longipetala</i> 055 Haemodoraceae	<i>Wurmbea pygmaea</i> 054J Colchicaceae
<i>Tribonanthes</i> <i>purpurea</i> 055 Haemodoraceae	<i>Wurmbea tenella</i> 054J Colchicaceae
<i>Tribulus</i> <i>terrestris</i> 173 Zygophyllaceae	<i>Xanthoparmelia digitiformis</i> L Parmeliaceae
<i>Trichodesma</i> <i>zeylanicum</i> var "unsorted" 310 Boraginaceae	<i>Xanthoparmelia hypoleia</i> L Parmeliaceae
<i>Tricoryne</i> <i>elatior</i> 054F Anthericaceae	<i>Xanthoparmelia isidiigera</i> L Parmeliaceae
<i>Trifolium</i> <i>dubium</i> 165 Papilionaceae	<i>Xanthoparmelia neotinctina</i> L Parmeliaceae
<i>Triglochin</i> aff <i>minutissimum</i> 026 Juncaginaceae	<i>Xanthoparmelia reptans</i> L Parmeliaceae
<i>Triglochin</i> <i>calcitratum</i> subsp <i>incurvum</i> ms 026 Juncaginaceae	<i>Xanthoparmelia substrigosa</i> L Parmeliaceae
<i>Triglochin</i> <i>centrocarpum</i> 026 Juncaginaceae	<i>Xanthoparmelia tasmanica</i> L Parmeliaceae
<i>Triglochin</i> <i>minutissimum</i> 026 Juncaginaceae	<i>Xanthoria parietina</i> L Teloschistaceae
<i>Triglochin</i> sp 026 Juncaginaceae	<i>Xanthosia candida</i> 281 Apiaceae
<i>Tripogon</i> <i>loliiformis</i> 031 Poaceae	<i>Xanthosia singuliflora</i> 281 Apiaceae
<i>Tripterococcus</i> <i>brunonis</i> 202 Stackhousiaceae	<i>Xerolirion divaricata</i> 054C Dasypogonaceae
<i>Triptilodiscus</i> <i>pygmaeus</i> 345 Asteraceae	