

# Opportunistic observation of predation of Loggerhead turtle hatchlings by feral cats on Dirk Hartog Island, Western Australia

S S Hilmer<sup>1</sup>, D Algar<sup>1</sup> & M Johnston<sup>2</sup>

<sup>1</sup>Department of Environment & Conservation, Science Division,  
PO Box 51, Wanneroo, WA, 6946.

<sup>2</sup>Department of Sustainability & Environment,  
Arthur Rylah Institute for Environmental Research,  
PO Box 137, Heidelberg, Vic., 3084.

*Manuscript received February 2010; accepted April 2010*

## Abstract

This note reports the opportunistic observation of predation of Loggerhead turtle hatchlings at Turtle Bay on Dirk Hartog Island, Western Australia. Data were collected on the movement patterns of three feral cats, fitted with GPS data-logger/radio-telemetry collars in the vicinity of turtle nests. This data as well as field observations demonstrate a distinct patrolling of Turtle Bay and surrounding areas. Furthermore, predation of turtle hatchlings by cats was confirmed by their presence in the stomach contents of one of the cats.

**Keywords:** feral cat, Loggerhead turtles, predation, Dirk Hartog Island

## Introduction

Globally, Loggerhead turtles (*Caretta caretta*) inhabit tropical, sub-tropical and temperate marine waters (Bolten and Witherington 2003). Within Australia, they are found in the eastern, northern and western waters of the continent (Prince 1994; Limpus 2008). Loggerhead turtles are listed as 'Endangered' under the Commonwealth *Environment Protection and Biodiversity Conservation Act* 1999 and the IUCN Red List (IUCN 2009).

Predation by feral cats is known to have a deleterious impact on endemic land vertebrates and breeding bird populations on both offshore and oceanic islands around the world (van Aarde 1980; Moors & Atkinson 1984; King 1985; Veitch 1985; Bloomer & Bester 1992; Bester *et al.* 2002; Keitt *et al.* 2002; Blackburn *et al.* 2004; Martinez-Gomez & Jacobsen 2004; Nogales *et al.* 2004). Predation by feral cats on Green turtle hatchlings (*Chelonia mydas*) on Aldabra Atoll, Seychelles has also been demonstrated (Seabrook 1989), although the impact could not be fully determined.

Turtle Bay and environs of Dirk Hartog Island is a major nesting ground for Loggerhead turtles with 800–1500 females breeding each year between November and March (DEWHA 2010). Footprints of feral cats around turtle nests were observed during a pilot study to assess the efficacy of a baiting program to eradicate feral cats on the island (Algar *et al.* submitted).

## Materials and Methods

In March 2009, sixteen feral cats were trapped on the island for a pilot study to assess bait efficacy. Two of these cats (DH27 and DH27.2) were caught near Turtle

Bay while another, MB8 was trapped close to the west coast (Figure 1). The cats were fitted with a GPS data-logger/radio-telemetry collars (small Feline/Possum GPS collar, weight: 105 g, dimensions: 55mm wide x 34 mm deep by 37 mm high under neck, Sirtrack Ltd, New Zealand), which collected and stored a location point every 40 minutes (DH27, DH27.2) and 10 minutes respectively (MB8). Cats were released at site of capture and not disturbed during the period of data collection.

A feral cat baiting program was conducted four weeks after the release of the cats (Johnston *et al.* 2009; Algar *et al.* submitted). Collars were retrieved following the death of individual cats, including DH27.2, following baiting. Cats DH27 and MB8 were shot at the conclusion of the study as they did not consume a toxic bait. No stomach analysis was conducted.

A 30 m section of beach to the west of Cape Inscription that encompassed an active turtle nest was inspected daily over 5 days for fresh feral cat activity. Observations and interpretations were made from the footprints made by cat(s), turtles, crabs and other species. Prints were cleared daily by sweeping the 'study area' with a broom.

## Results

Each of these cats was male and weighed; DH27, 5.1 kg, DH27.2, 4.5 kg and MB8, 5.5 kg when trapped. A compilation of the location data derived from GPS data-logger collars, between 30 March and 30 April 2009, is shown in Figure 2. The data indicate almost linear walking patterns as the cats foraged along the beach. In addition, fresh cat tracks, thought to be MB8, were seen daily at the turtle nesting sites as shown in Figure 3. Toxicosis following ingestion of the poison '1080' (sodium monofluoroacetate) used in the feral cat baiting program often results in regurgitation of stomach contents. Vomitus of an entire turtle hatchling and





Figure 1. Location map of study area, Turtle Bay on Dirk Hartog Island (circled area).



**Figure 2.** GPS-locations of three feral cats at and near Turtle Bay collected over a period of 10 days (MB8, 29.3-8.4.2009) and 30 days (DH27, DH27.2, 30.3- 30.4.2009).



Figure 3. Cat tracks on Turtle Bay beach next to Loggerhead Turtle nests (left from cat tracks).

unidentified rodent was located within 10 metres of the carcass of DH27.2 (see Figure 4).

### Discussion

Many terrestrial species, both native and introduced, are known to predate turtle eggs or emergent hatchlings (Limpus 1973; Environment Australia 2003; Limpus 2008) however, predation by feral cats has rarely been demonstrated (Seabrook 1989). Hamann *et al.* (2006) suggested that predation by feral cats on Flatback turtles (*Natator depressus*) may be of management concern on West Island in the Sir Edward Pellew group, Northern Territory however, despite the presence of many cat

tracks at nest sites, predation by cats was not observed. The finding reported in this note, despite a small data set, confirms that feral cats, known to be opportunistic hunters (Jones & Coman 1981; Dickman 1996; Paltridge *et al.* 1997; Risbey *et al.* 1999), predate Loggerhead turtle hatchlings and provides further confirmation that feral cats will consume large prey items (Marks *et al.* 2006; Hetherington *et al.* 2007). The regurgitated prey items, (Figure 4) were entire and stained with the Rhodamine B dye used in the poison baits. The extent of the predation by feral cats on Dirk Hartog Island on turtle hatchlings is not known, but a planned feral cat eradication program on the island, proposed to commence in 2011 may improve hatching survival of Loggerhead turtles at Turtle Bay.



**Figure 4.** Regurgitated Loggerhead Turtle hatchling found near cat DH27. 2 (Note: red dye is Rhodamine B that was incorporated into the baits).

**Acknowledgments:** Additional field assistance was provided by Mike Onus, Neil Hamilton, Bruce Withnell, Katrin Koch, David Hawke and Steve Virgin of the WA Department of Conservation and Environment (DEC). Logistic support was provided by the staff at the DEC Denham office especially Brett Fitzgerald. Advice regarding Loggerhead turtle biology was sourced from Bob Prince, Linda Reinhold and Sabrina Trocini.

## References

- Algar D, Johnston M, Hilmer S (submitted) A pilot study for the proposed eradication of feral cats on Dirk Hartog Island, Western Australia. *In: Island Invasives: Eradication and Management*. Auckland.
- Bester M N, Bloomer J P, van Aarde R J, Erasmus B H, van Rensburg P J J, Skinner J D, Howell P G & Naude T W 2002 A review of the successful eradication of feral cats from sub-Antarctic Marion Island, Southern Indian Ocean. *South African Journal Wildlife Research* 32: 65–73.
- Blackburn T M, Cassey P, Duncan R P, Evans K L & Gaston K J 2004 Avian extinction and mammalian introductions on oceanic islands. *Science* 305: 1955–1958.
- Bloomer J P & Bester M N 1992 Control of feral cats on sub-Antarctic Marion Island, Indian Ocean. *Conservation Biology* 60: 211–219.
- Bolten A & Witherington B 2003 *Loggerhead Sea Turtles*. Smithsonian Books, Washington.
- Department of the Environment, Water, Heritage & the Arts 2010 *Caretta caretta* in Species Profile and Threats Database, Department of the Environment, Water, Heritage & the Arts, Canberra.
- [http://www.environment.gov.au/cgi-bin/sprat/public/publicspecies.pl?taxon\\_id=1763](http://www.environment.gov.au/cgi-bin/sprat/public/publicspecies.pl?taxon_id=1763) accessed January 27, 2010.
- Dickman C R 1996 Overview of the impact of feral cats on Australian native fauna. Report to Australian Nature Conservation Agency.
- Environment Australia 2003 Recovery plan for marine turtles in Australia. Commonwealth of Australia (ISBN 0642214360).
- Hamann M, Schauble C, Simon T, Johnson J, Evans S, Dorr T & Kennett R 2006 Sea turtles nesting in the Sir Edward Pellew Islands, Gulf of Carpentaria, Northern Territory. *Memoirs of the Queensland Museum*, 52 (1): 71–78.
- Hetherington C, Algar D, Mills H R & Bencini R 2007 Increasing the target-specificity of ERADICAT® for feral cat (*Felis catus*) control by encapsulating a toxicant. *Wildlife Research* 34: 1–5.
- IUCN 2009. IUCN Red List of Threatened Species. Version 2009.2. <[www.iucnredlist.org](http://www.iucnredlist.org)>.
- Johnston M, Algar D, Onus M, Hamilton N, Hilmer S, Withnell B & Koch K 2009 A bait efficacy trial for the management of feral cats on Dirk Hartog Island. Arthur Rylah Institute for Environmental Research Client Report. Department of Sustainability & Environment, Heidelberg, Victoria.

- Jones E & Coman B J 1981 Ecology of the feral cat, *Felis catus* (L.), in South-Eastern Australia I. Diet. *Wildlife Research* 8 (3): 537–547.
- Keitt B S, Wilcox C, Tershy B R, Croll D A & Donlan C J 2002 The effect of feral cats on the population viability of blackvented shearwaters (*Puffinus opisthomelas*) on Natividad Island, Mexico. *Animal Conservation* 5: 217–223.
- King W B 1985 Island birds: will the future repeat the past? *In: Conservation of island birds* (ed P J Moors), ICBP Technical Publication No. 3, 3–15.
- Limpus C J 1973 Avian predators of sea turtles in south-east Queensland rookeries. *The Sunbird* 4: 45–51.
- Limpus C J 2008 A biological review of Australian Marine Turtles. 1. Loggerhead Turtle *Caretta caretta* (Linnaeus). Queensland Environment Protection Agency.
- Marks C A, Johnston M J, Fisher P M, Pontin K & Shaw M J 2006 Differential particle size: promoting target-specific baiting of feral cats. *Journal of Wildlife Management* 70: 1119–1124.
- Martinez-Gomez J E & Jacobsen J K 2004 The conservation status of Townsend's shearwater *Puffinus auricularis auricularis*. *Conservation Biology* 116: 35–47.
- Moors P J & Atkinson I A E 1984 Predation on seabirds by introduced animals, and factors affecting its severity. *In: Status and conservation of the world's seabirds* (eds J P Croxall, P J H Evans & R W Schreiber). ICBP Technical Publication No. 2, 667–690.
- Nogales M, Martin A, Tershy B R, Donlan C J, Veitch D, Puerta N, Wood B & Alonso, J 2004 A review of feral cat eradication on islands. *Conservation Biology* 18: 310–319.
- Paltridge R, Gibson D & Edwards G 1997 Diet of the feral cat (*Felis catus*) in central Australia. *Wildlife Research* 24: 67–76
- Prince R I 1994 Status of the Western Australian marine turtle populations: the Western Australian Marine Turtle Project 1986–1990. *In: Proceedings of the Australian Marine Turtle Conservation Workshop, Gold Coast 14–17 November 1990* (ed J Russell) Queensland Department of Environment and Heritage. Canberra, ANCA, 1–14.
- Risbey D A, Calver M C & Short J 1999 The impact of cats and foxes on the small vertebrate fauna of Heirisson Prong, Western Australia. I. Exploring potential impact using diet analysis. *Wildlife Research* 26: 621–630.
- Seabrook W 1989 Feral cats (*Felis catus*) as predators of hatchling green turtles (*Chelonia mydas*). *Journal of Zoology* 219: 83–88.
- van Aarde R J 1980 The diet and feeding behaviour of feral cats, *Felis catus*, on Marion Island. *South African Journal of Wildlife Research* 10: 123–128.
- Veitch C R 1985 Methods of eradicating feral cats from offshore islands in New Zealand. *In: Conservation of island birds* (P J Moors ed.). ICBP Technical Publication No. 3, 125–141